



# **Performance of *Beauveria bassiana* (Balsamo) Vuillemin Stored as Oil Cultures under Different Temperature Regimes**

**Chelsia R<sup>a++\*</sup>, Manjula K<sup>a#</sup>, Harathi P N<sup>b†</sup>,  
Gurivi Reddy M<sup>c‡</sup> and Abdul Mujeeb S<sup>a^</sup>**

<sup>a</sup> Department of Entomology, S.V. Agricultural College, Tirupati-517 502, (Acharya N.G. Ranga Agricultural University, Andhra Pradesh), India.

<sup>b</sup> Department of Entomology, Regional Agricultural Research Station, Tirupati-517 502, Andhra Pradesh, India.

<sup>c</sup> Department of Plant Pathology, Agricultural College, Mahanandi- 518 502, Andhra Pradesh, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/mrji/2024/v34i81470>

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc. are available here: <https://www.sdiarticle5.com/review-history/120760>

**Original Research Article**

**Received: 23/05/2024**

**Accepted: 25/07/2024**

**Published: 29/07/2024**

<sup>++</sup> M. Sc. Scholar;

<sup>#</sup> Professor and Head;

<sup>†</sup> Scientist;

<sup>‡</sup> Assistant Professor;

<sup>^</sup> Ph.D. Scholar;

\*Corresponding author: E-mail: [rchelsia10@gmail.com](mailto:rchelsia10@gmail.com);

**Cite as:** R, Chelsia, Manjula K, Harathi P N, Gurivi Reddy M, and Abdul Mujeeb S. 2024. "Performance of *Beauveria Bassiana* (Balsamo) Vuillemin Stored As Oil Cultures under Different Temperature Regimes". *Microbiology Research Journal International* 34 (8):45-58. <https://doi.org/10.9734/mrji/2024/v34i81470>.

## ABSTRACT

Entomopathogenic Fungal agents play an important role in controlling pest population. Besides isolating and exploiting the virulent fungal agents it is essential to maintain its viability for longer period at affordable cost, thereby they could be employed for further research. Usage of oils and storage at lower temperatures prevent conidial desiccations and could be comparatively a cheaper alternative to freeze drying and cryopreservation to maintain its virulence for extended periods. Oil cultures of *Beauveria bassiana* (Balsamo) Vuillemin were prepared from the laboratory-maintained isolate by using two vegetable oils viz., Rice Bran oil, Sesamum oil and two mineral oils viz., Liquid Paraffin oil and Heavy grade Mineral Oil. Prepared oil cultures were stored under three different temperature regimes viz., at room temperature ( $28 \pm 3^\circ\text{C}$ ), refrigerated condition ( $4^\circ\text{C}$ ) and freezing temperature ( $-20^\circ\text{C}$ ). These 12 cultures stored at different temperatures along with crude culture and untreated control were evaluated for viability and virulence against third instar *Spodoptera litura* (Fabricius) larvae on the day of preparation (0 DAS) and upto 180 days under laboratory conditions. Among all oil cultures, Sesamum oil cultures and Liquid Paraffin oil cultures stored at  $-20^\circ\text{C}$  performed well with highest conidial germination of 87.85% and 83.57% and highest larval mortality of 90.00% and 86.67% on the day of preparation (0 DAS). After 180 DAS of storage, conidial germination in these two formulations was in the range of 62.87% to 61.43% and larval mortality was 60.00%. Comparatively, least performance was observed in mineral oil formulations stored at room temperature ( $T_1$ ) and in unformulated crude cultures.

**Keywords:** *Beauveria bassiana*; oil cultures; vegetable oils; mineral oils; different temperature regimes; storage; viability; virulence.

## 1. INTRODUCTION

According to FAO's 2022 report, India consumed over 61,000 tonnes (t) of pesticides in 2020 [1]. These pesticides in addition to creating pesticide residues, environmental pollution issues and other issues, the widespread use of synthetic pesticides inevitably lead to the development of insecticide resistance, harmful effect on natural enemies, insecticide-induced resurgence of the pests and biodiversity decline [2-4]. Considering these limitations, environmentally benign pest management methods have become increasingly popular to rebuild the balance of cropland ecosystems for the sustainable control of crop pests [5]. Pest management by biocontrol agents are assuming prominence and have been considered as an important strategy in insect population reduction [6]. Among biocontrol agents, microbial pathogens such as bacteria, fungi, viruses, nematodes and protozoans are promising agents for the effective control of insects.

Entomopathogenic fungi are regarded as the most significant among the numerous microbial agents due to their wide host range, eco-friendly nature, effectiveness against target pest etc. They are cheaper in long run, show lesser residual effects, and are able to overcome the problem of resistance [7]. EPF is advantageous in pest control because direct ingestion of fungal

propagules is not needed by insects, thus also becoming active against the non-feeding stages of insects [8]. They cause insect mortality by nutritional deficiency, destruction of tissues and by release of toxins. The entry of entomopathogenic fungi through the insect cuticle occur by a combination of mechanical pressure and enzymatic degradation [9]. They have the ability to synthesize extracellular cuticle-degrading enzymes like chitinase, protease and lipase that work together to overcome the chitinous and proteinaceous components of the insect cuticle, penetrating the germ tube inside thus, enabling hyphal access to the haemolymph and colonizing the insects to death [10-12].

Several species of entomopathogenic fungi have now been industrialized [13]. *Beauveria bassiana* is the asexual form (anamorph) of *Cordyceps bassiana* fungus that infects a huge variety of insects, which is used to control crop infestations caused by aphids, thrips and whiteflies [14] as well as lepidopteran pests [15]. Among the cyclic hexadepsipeptide mycotoxins produced by the different EPF, beauvericin produced by them have shown the most effective larvicidal properties [16]. Even though, effective strains were isolated there was still problem with the development of suitable cost-effective formulations which could overcome harsh environmental conditions and ensure biological

and chemical stability as well as viability during storage for extended periods [17].

One among the technique is the usage of oils for the preservation of entomopathogenic fungal formulations. It has been suggested that oil formulations can prevent conidial desiccation and improve adhesion of conidia to the hydrophobic surface of insect cuticle and enabling the opportunistic spread of conidia in to high humidity locations such as the inter-segmental membranes [18-20]. In field conditions, higher temperature, lower humidity and Ultraviolet radiation rays (UV) exposure pose detrimental effects on fungal conidia. This situation warrants shift to the use of oil based formulations which showed good control of insect pests under field condition [21]. Oil affords protection to fungal conidia from the UV rays of sunlight thereby improvement in the field performance [22].

While, cryoprotectant preservation at  $-196^{\circ}\text{C}$  submerged in liquid nitrogen following freezing at a controlled rate is generally accepted as the optimal storage technique for preserving both cell viability and biological properties *i.e.* infectivity and virulence, it is considerably expensive and time consuming which is not affordable by many laboratories [23,24]. Maintenance at optimum temperature for long term preservation of EPF culture is of utmost importance.

Therefore, its necessary for the development of suitable and affordable formulation technologies for the proper maintenance and preservation of different entomopathogenic fungal cultures. There have been relatively fewer studies conducted on optimising the storage conditions for *Beauveria bassiana* oil cultures, which could remain viable for extended periods and be suitable for later retrieval. Therefore, the present study was undertaken to evaluate the performance of *B. bassiana* stored as oil cultures under different temperature regimes.

## 2. MATERIALS AND METHODS

**Place and duration of study:** Insect Pathology laboratory, Central Instrumentation Laboratory and Insectary, Department of Entomology Sri Venkateswara Agricultural College, Tirupati during between September, 2023 and April, 2024.

### 2.1 Preparation of *Beauveria bassiana* Oil Cultures

For preparation of oil cultures, 4 oils were chosen *viz.*, 2 vegetable oils (Rice bran oil,

Sesamum oil) and 2 mineral oils (Heavy grade Mineral oil, Liquid paraffin oil). Saboraud's Dextrose Agar with Yeast Extract Medium (SDAY) was prepared, poured into test tubes, autoclaved at  $121^{\circ}\text{C}$  at 15 Psi for 15 minutes and made as slants. After slants got cooled down, discs or loop of spores and mycelia of laboratory-maintained culture of *B. bassiana* were inoculated into slants under aseptic condition and incubated at  $25 \pm 2^{\circ}\text{C}$  temperature. Ten to twelve days after inoculation, satisfactory growth and sporulation of *B. bassiana* were obtained in almost all slants (Fig. 1). Each fungal slant was poured with autoclaved vegetable and mineral oils *viz.*, Rice bran oil, Sesamum oil, Mineral oil and Liquid paraffin oil respectively until the entire fungi grown in slant media got submerged with oil (Fig. 2).



Fig. 1. Slant culture of *B. bassiana*



Fig. 2. *B. bassiana* oil cultures

Humber [25] reported that storage of culture slants under a layer of sterile mineral oil was one of the oldest, simplest, and least expensive methods for long-term culture preservation.

## 2.2 Storage of *Beauveria bassiana* oil Cultures at Different Temperature Regimes

The prepared oil cultures of *B. bassiana* with Rice bran oil, Sesamum oil, Mineral oil and Liquid paraffin oil were maintained at room temperature ( $28 \pm 3^\circ\text{C}$ ), refrigerated condition ( $4^\circ\text{C}$ ) in Refrigerator at Insect Pathology laboratory and freezing temperature ( $-20^\circ\text{C}$ ) in Deep Freezer at Central instrumentation Lab of the college.

A total of treatments 14 treatments (including control) of *B. bassiana* cultures were prepared.

- T<sub>1</sub> : Mineral oil culture of *B. bassiana* stored at Room temperature ( $28 \pm 3^\circ\text{C}$ )
- T<sub>2</sub> : Mineral oil culture of *B. bassiana* stored at Refrigerated temperature ( $4^\circ\text{C}$ )
- T<sub>3</sub> : Mineral oil culture of *B. bassiana* stored at freezing temperature ( $-20^\circ\text{C}$ )
- T<sub>4</sub> : Liquid paraffin oil culture of *B. bassiana* stored at Room temperature ( $28 \pm 3^\circ\text{C}$ )
- T<sub>5</sub> : Liquid paraffin oil culture of *B. bassiana* stored at Refrigerated temperature ( $4^\circ\text{C}$ )
- T<sub>6</sub> : Liquid paraffin oil culture of *B. bassiana* stored at freezing temperature ( $-20^\circ\text{C}$ )
- T<sub>7</sub> : Rice bran oil culture of *B. bassiana* stored at Room temperature ( $28 \pm 3^\circ\text{C}$ )
- T<sub>8</sub> : Rice bran oil culture of *B. bassiana* stored at Refrigerated temperature ( $4^\circ\text{C}$ )
- T<sub>9</sub> : Rice bran oil culture of *B. bassiana* stored at freezing temperature ( $-20^\circ\text{C}$ )
- T<sub>10</sub> : Sesamum oil culture of *B. bassiana* stored at Room temperature ( $28 \pm 3^\circ\text{C}$ )
- T<sub>11</sub> : Sesamum oil culture of *B. bassiana* stored at Refrigerated temperature ( $4^\circ\text{C}$ )
- T<sub>12</sub> : Sesamum oil culture of *B. bassiana* stored at freezing temperature ( $-20^\circ\text{C}$ )
- T<sub>13</sub> : Unformulated culture of *B. bassiana* (Not preserved in oil)
- T<sub>14</sub> : Untreated control

Murugasridevi et al. [26] prepared oil-based formulations of *Beauveria bassiana* in 100 ml of paraffin oil, along with adjuvants like polyethylene glycol and tween 80 and stored it under different storage conditions viz. ambient temperature ( $28 \pm 3^\circ\text{C}$ ) and refrigerated condition ( $4^\circ\text{C}$ ) and tested its performance for 24 weeks.

## 2.3 Evaluation of *B. bassiana* oil Cultures Stored under Different Temperature Regimes

The viability of *B. bassiana* conidia and its efficacy was studied on the day of preparation (0 DAS) and at monthly intervals up to 180 days (DAS).

### 2.3.1 Assessment of viability of oil cultures

Serial dilutions of  $1 \times 10^5$  spores  $\text{ml}^{-1}$  of all oil cultures were employed for viability experiments. Spore count was taken with the help of haemocytometer. Two to three drops of spore suspension were placed in the cavity slide. The cavity slide was prepared by arranging moistened cotton in petriplates and incubated in the incubator at  $22^\circ\text{C}$ . After 24 hrs, the spore suspension was observed under microscope for counting total number of spores and number of germinated spores. The germination percentage of *Beauveria bassiana* conidia at different treatments were calculated.

### 2.3.2 Assessment of virulence of oil cultures against third instar *Spodoptera litura* larvae

From each treatment, 0.5 ml of oil culture was poured into 100 ml of distilled water and added with 100  $\mu\text{l}$  of Triton- X 100 (0.1 %) into conical flasks to make spray suspension. Triton- X 100 (0.1 %) as a wetting agent for uniform dispersion of spores with oil. Spore suspensions of  $1 \times 10^7$  spores per ml were standardized after assessing the number of spores in the suspension with an improved Neubauer haemocytometer. Spray suspensions were sprayed on castor leaves with atomizer and air dried. Freshly moulted third instar larvae were allowed to feed on them. The treatments were replicated thrice to confirm the reproducibility of the results. The bioassays were conducted at room temperature under laboratory conditions. From next day, larvae were fed with fresh castor leaves. Daily observations on post treatment changes in larvae and larval mortalities were recorded. Sandhu et al. [27] evaluated the effects of temperature ( $0^\circ$ ,  $10^\circ$ ,  $20^\circ$ ,  $30^\circ$  and  $40^\circ\text{C}$ ) on conidial viability and virulence of *Beauveria bassiana* against third instar *Helicoverpa armigera* (Hubner) larvae over a 24-month period.

## 2.4 Analysis of the Data

All the recorded observations i.e. conidial viability and larval mortality were converted to

percentage values. The observations were statistically analysed by using SPSS v 16 software. Data were subjected to analysis of variance (ANOVA) at  $P < 0.01$  level of significance. Means were compared by Duncan's Multiple Range Test (DMRT) [28].

The conidial germination was expressed as per cent conidial viability by using the formula:

$$\text{Per cent conidia viability} = \frac{\text{Number of conidia germinated}}{\text{Total conidia}} \times 100$$

The larval mortality was expressed as per cent larval mortality by using the formula:

$$\text{Per cent larval mortality} = \frac{\text{Number of larvae dead due to infection}}{\text{Total number of larvae treated}} \times 100$$

### 3. RESULTS AND DISCUSSION

A total of twelve types of oil cultures were prepared by using four oils and these cultures were maintained at three different temperatures. Additionally, an unformulated crude culture of *Beauveria bassiana* and untreated control were maintained. The post treatment larval changes and the results of viability and virulence were presented below.

After releasing the larvae onto treated leaves, it was observed that there was a noticeable reduction in their tendency to feed, particularly on leaves treated with sesame oil. When the larvae were fed with fresh leaves the day after treatment, they exhibited a reduced inclination to feed. The room temperature ( $28 \pm 3^\circ\text{C}$ ) was highly conducive for mycosis development. Freshly moulted early 3<sup>rd</sup> instar larvae were significantly more susceptible to EPF treatments compared to late 3<sup>rd</sup> and 4<sup>th</sup> instar *S. litura* larvae. After 2-3 days of treatment with *B. bassiana*, the infected larvae exhibited sluggish movement and consumed less food material. The larval integument got shrunken and larval bodies had become exceedingly smooth. Majority of larvae were dead within 6 days of inoculation with fungal spores. There was sparse growth of fungal mycelium on the surface of larval integument and whitish mycelial growth of fungus was conspicuous from 5<sup>th</sup> day after inoculation. The entire larval body was covered with puffy whitish mycelia and whitish fungal spores was produced on to larval surface. As the time progressed, the puffiness of the fungus on the larval body reduced gradually. The larval body eventually turned into a hard and mummified cadaver. The stiffness of the cadaver after death

might be due to excessive fungal growth inside larval body. Larval mortality occurred more rapidly within 5-6 days with oil formulations, compared to 8-10 days with unformulated crude suspensions. The results were in agreement with Ummidi and Vadlamani [6] who conducted bioassay studies of oil formulations of *B. bassiana* against *S. litura*. Based on compatibility studies, almond oil, olive oil, gingelly oil, castor oil based formulations were chosen for bioassay. He reported that all the four formulations displayed higher mortalities of the target pest compared to unformulated conidia, with *B. bassiana* showed higher mortality with gingelly oil and almond oil.

In the present study it was observed that, the infected larvae progressed to next instar and pupated earlier compared to untreated larvae *i.e.* significant decrease in larval period was observed due to infection in comparison with untreated control. Some affected larvae metamorphosed to malformed pupae and adult. These results corroborate with the findings of Torrado-Leon *et al.* [29] who documented the interference in the moulting process of *Bemisia tabacci* (Gennadius) nymphs when treated with *B. bassiana*. More than 30% of the imagos emerged from treated nymphs were unable to detach completely from the exuvium.

It was noticed that on the day of preparation, the highest mean per cent conidial viability of 87.85% was recorded in Sesamum oil cultures (T<sub>10</sub>, T<sub>11</sub>, T<sub>12</sub>), followed by Liquid paraffin oil cultures (T<sub>4</sub>, T<sub>5</sub>, T<sub>6</sub>) and Rice bran oil cultures (T<sub>7</sub>, T<sub>8</sub>, T<sub>9</sub>) which recorded 83.57% and 80.71% respectively (Fig. 3). Comparatively lower conidial viability of 77.14% was observed in oil cultures formulated using Mineral oil (T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>). The unformulated culture (T<sub>13</sub>) recorded a conidial viability of 74.28% (Table. 1). The data regarding the mean per cent larval mortality indicated that the maximum larval mortality of 90.00% was recorded in Sesamum oil cultures (T<sub>10</sub>, T<sub>11</sub>, T<sub>12</sub>), followed by Liquid paraffin oil cultures (T<sub>4</sub>, T<sub>5</sub>, T<sub>6</sub>) and Rice bran oil cultures (T<sub>7</sub>, T<sub>8</sub>, T<sub>9</sub>) which recorded 86.67% and 83.33% respectively (Fig. 4). Comparatively lower larval mortality of 80.00% was observed in oil cultures formulated using Mineral oil (T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>). The unformulated culture (T<sub>13</sub>) recorded a larval mortality of 76.67%. There was no larval mortality recorded in untreated control T<sub>14</sub>. (Table. 2).

**Table 1. Viability of *Beauveria bassiana* conidia stored under different temperature regimes after formulating as oil cultures (2023-2024)**

S. No.	Treatment	Mean Per cent Conidia germination						
		0 DAS	30 DAS	60 DAS	90 DAS	120 DAS	150 DAS	180 DAS
1	T <sub>1</sub>	77.14 <sup>c</sup> (61.44)	67.85 <sup>cd</sup> (55.46)	61.43 <sup>de</sup> (51.61)	57.87 <sup>d</sup> (49.53)	51.43 <sup>cd</sup> (45.82)	44.29 <sup>de</sup> (41.72)	41.43 <sup>f</sup> (40.07)
2	T <sub>2</sub>	77.14 <sup>c</sup> (61.44)	70.71 <sup>bcd</sup> (57.23)	65.71 <sup>cde</sup> (54.16)	62.87 <sup>cd</sup> (52.46)	58.57 <sup>bcd</sup> (49.93)	51.43 <sup>bcd</sup> (45.82)	48.57 <sup>def</sup> (44.18)
3	T <sub>3</sub>	77.14 <sup>c</sup> (61.44)	74.28 <sup>abc</sup> (59.53)	70.71 <sup>bcd</sup> (57.23)	65.00 <sup>bc</sup> (53.73)	64.28 <sup>abc</sup> (53.30)	58.57 <sup>abcd</sup> (49.93)	54.29 <sup>bcd</sup> (47.46)
4	T <sub>4</sub>	83.57 <sup>b</sup> (66.09)	70.71 <sup>bcd</sup> (57.23)	67.85 <sup>bcd</sup> (55.46)	61.43 <sup>cd</sup> (51.61)	54.29 <sup>bcd</sup> (47.46)	49.29 <sup>bcd</sup> (44.59)	45.71 <sup>def</sup> (42.54)
5	T <sub>5</sub>	83.57 <sup>b</sup> (66.09)	74.28 <sup>abc</sup> (59.53)	70.71 <sup>bcd</sup> (57.23)	65.00 <sup>bc</sup> (53.73)	61.43 <sup>abc</sup> (51.61)	58.57 <sup>abcd</sup> (49.93)	51.43 <sup>cde</sup> (45.82)
6	T <sub>6</sub>	83.57 <sup>b</sup> (66.09)	79.28 <sup>ab</sup> (62.92)	74.28 <sup>abc</sup> (59.53)	70.71 <sup>ab</sup> (57.23)	65.71 <sup>abc</sup> (54.16)	63.75 <sup>abc</sup> (52.98)	61.43 <sup>ab</sup> (51.61)
7	T <sub>7</sub>	80.71 <sup>bc</sup> (63.95)	67.85 <sup>cd</sup> (55.46)	64.28 <sup>de</sup> (53.30)	57.87 <sup>d</sup> (49.53)	54.29 <sup>bcd</sup> (47.46)	48.57 <sup>cde</sup> (44.18)	44.29 <sup>ef</sup> (41.72)
8	T <sub>8</sub>	80.71 <sup>bc</sup> (63.95)	75.71 <sup>abc</sup> (60.47)	69.28 <sup>bcd</sup> (56.34)	65.71 <sup>bc</sup> (54.16)	58.57 <sup>bcd</sup> (49.93)	55.71 <sup>abcd</sup> (48.28)	51.43 <sup>cde</sup> (45.82)
9	T <sub>9</sub>	80.71 <sup>bc</sup> (63.95)	76.42 <sup>abc</sup> (60.95)	75.71 <sup>ab</sup> (60.47)	73.57 <sup>a</sup> (59.06)	69.28 <sup>ab</sup> (56.34)	64.28 <sup>ab</sup> (53.30)	58.57 <sup>abc</sup> (49.93)
10	T <sub>10</sub>	87.85 <sup>a</sup> (69.60)	70.71 <sup>bcd</sup> (57.23)	65.17 <sup>de</sup> (53.83)	63.57 <sup>cd</sup> (52.87)	57.87 <sup>bcd</sup> (49.53)	49.29 <sup>bcd</sup> (44.59)	47.85 <sup>def</sup> (43.77)
11	T <sub>11</sub>	87.85 <sup>a</sup> (69.60)	80.71 <sup>ab</sup> (63.95)	76.42 <sup>ab</sup> (60.95)	74.28 <sup>a</sup> (59.53)	67.85 <sup>ab</sup> (55.46)	62.87 <sup>abc</sup> (52.46)	54.29 <sup>bcd</sup> (47.46)
12	T <sub>12</sub>	87.85 <sup>a</sup> (69.60)	82.14 <sup>a</sup> (65.00)	79.28 <sup>a</sup> (62.92)	75.71 <sup>a</sup> (60.47)	74.28 <sup>a</sup> (59.53)	67.85 <sup>a</sup> (55.46)	62.87 <sup>a</sup> (52.46)
13	T <sub>13</sub>	74.28 <sup>d</sup> (52.87)	61.43 <sup>d</sup> (51.61)	58.57 <sup>e</sup> (49.93)	51.43 <sup>e</sup> (45.82)	45.71 <sup>d</sup> (42.54)	39.29 <sup>e</sup> (38.82)	32.85 <sup>g</sup> (34.97)
14	T <sub>14</sub>	0.00 <sup>e</sup> (0.00)	0.00 <sup>e</sup> (0.00)	0.00 <sup>f</sup> (0.00)	0.00 <sup>f</sup> (0.00)	0.00 <sup>f</sup> (0.00)	0.00 <sup>f</sup> (0.00)	0.00 <sup>h</sup> (0.00)
	General mean	75.10	68.01	64.24	60.36	55.97	50.98	46.79
	F	Sig.	Sig.	Sig.	Sig.	Sig.	Sig.	Sig.
	Sig.	0.000	0.000	0.000	0.000	0.000	0.000	0.000
	SE(m)±	0.72	0.37	0.61	0.59	0.57	0.56	0.56
	C.D.	2.10	1.07	1.78	1.72	1.67	1.64	1.63
	C.V.	2.09	1.16	2.02	2.05	2.10	2.20	2.31

DAS: Days After Storage, Values are the means of three replications,

Values in parentheses are angular transformed values

Means in a column followed by same superscript are not significantly different according to DMRT at P≤ 0.05.

SE(m)± = Standard error of mean C.V. = Coefficient of variation

C.D. = Critical Difference at 1% level of significance

T<sub>1</sub>: Mineral oil culture of *B. bassiana* stored at Room temperature (28 ± 3°C), T<sub>2</sub>:Mineral oil culture of *B. bassiana* stored at Refrigerated temperature (4°C), T<sub>3</sub>:Mineral oil culture of *B. bassiana* stored at freezing temperature (-20°C), T<sub>4</sub>:Liquid paraffin oil culture of *B. bassiana* stored at Room temperature (28 ± 3°C), T<sub>5</sub>:Liquid paraffin oil culture of *B. bassiana* stored at Refrigerated temperature (4°C), T<sub>6</sub>:Liquid paraffin oil culture of *B. bassiana* stored at freezing temperature (-20°C), T<sub>7</sub>:Rice bran oil culture of *B. bassiana* stored at Room temperature (28 ± 3°C), T<sub>8</sub>:Rice bran oil culture of *B. bassiana* stored at Refrigerated temperature (4°C), T<sub>9</sub>:Rice bran oil culture of *B. bassiana* stored at freezing temperature (-20°C), T<sub>10</sub>:Sesamum oil culture of *B. bassiana* stored at Room temperature (28 ± 3°C), T<sub>11</sub>:Sesamum oil culture of *B. bassiana* stored at Refrigerated temperature (4°C), T<sub>12</sub>:Sesamum oil culture of *B. bassiana* stored at freezing temperature (-20°C), T<sub>13</sub>:Unformulated culture of *B. bassiana* (Not preserved in oil), T<sub>14</sub>:Untreated control.

**Table 2. Virulence of *Beauveria bassiana* conidia stored under different temperature regimes after formulating as oil cultures against third instar larva of *Spodoptera litura* under laboratory conditions (2023-2024)**

S. No	Treatment	Mean per cent larval Mortality of third instar <i>Spodoptera litura</i> larvae						
		0 DAS	30 DAS	60 DAS	90 DAS	120 DAS	150 DAS	180 DAS
1	T <sub>1</sub>	80.00 <sup>a</sup> (63.44)	66.67 <sup>de</sup> (54.74)	60.00 <sup>de</sup> (50.77)	56.67 <sup>cd</sup> (48.33)	50.00 <sup>de</sup> (45.00)	43.33 <sup>cd</sup> (41.17)	40.00 <sup>cd</sup> (39.23)
2	T <sub>2</sub>	80.00 <sup>a</sup> (63.44)	70.00 <sup>cde</sup> (56.79)	66.67 <sup>cde</sup> (54.74)	60.00 <sup>bcd</sup> (50.77)	56.67 <sup>bcde</sup> (48.83)	50.00 <sup>abcd</sup> (45.00)	46.67 <sup>abc</sup> (43.09)
3	T <sub>3</sub>	80.00 <sup>a</sup> (63.44)	76.67 <sup>bcd</sup> (61.12)	70.00 <sup>bcd</sup> (56.79)	66.67 <sup>abc</sup> (54.74)	63.33 <sup>abcd</sup> (52.73)	56.67 <sup>abc</sup> (48.83)	53.33 <sup>abc</sup> (46.91)
4	T <sub>4</sub>	86.67 <sup>a</sup> (68.59)	70.00 <sup>cde</sup> (56.79)	66.67 <sup>cde</sup> (54.74)	60.00 <sup>bcd</sup> (50.77)	53.33 <sup>cde</sup> (46.91)	50.00 <sup>abcd</sup> (45.00)	43.33 <sup>bcd</sup> (41.17)
5	T <sub>5</sub>	86.67 <sup>a</sup> (68.59)	73.33 <sup>bcd</sup> (58.91)	70.00 <sup>bcd</sup> (56.79)	66.67 <sup>abc</sup> (54.74)	60.00 <sup>abcd</sup> (50.77)	56.67 <sup>abc</sup> (48.83)	50.00 <sup>abc</sup> (45.00)
6	T <sub>6</sub>	86.67 <sup>a</sup> (68.59)	83.33 <sup>ab</sup> (65.90)	76.67 <sup>abc</sup> (61.12)	70.00 <sup>abc</sup> (56.79)	66.67 <sup>abc</sup> (54.74)	63.33 <sup>ab</sup> (52.73)	60.00 <sup>a</sup> (50.77)
7	T <sub>7</sub>	83.33 <sup>a</sup> (65.90)	66.67 <sup>de</sup> (54.74)	63.33 <sup>de</sup> (52.73)	56.67 <sup>cd</sup> (48.83)	53.33 <sup>cde</sup> (46.91)	46.67 <sup>bcd</sup> (43.09)	43.33 <sup>bcd</sup> (41.17)
8	T <sub>8</sub>	83.33 <sup>a</sup> (65.90)	76.67 <sup>bcd</sup> (61.12)	70.00 <sup>bcd</sup> (56.79)	63.33 <sup>bcd</sup> (52.73)	56.67 <sup>bcde</sup> (48.83)	53.33 <sup>abcd</sup> (46.91)	50.00 <sup>abc</sup> (45.00)
9	T <sub>9</sub>	83.33 <sup>a</sup> (65.90)	80.00 <sup>abc</sup> (63.44)	76.67 <sup>abc</sup> (61.12)	73.33 <sup>ab</sup> (58.91)	70.00 <sup>ab</sup> (56.79)	63.33 <sup>ab</sup> (52.73)	56.67 <sup>ab</sup> (48.83)
10	T <sub>10</sub>	90.00 <sup>a</sup> (71.57)	70.00 <sup>cde</sup> (56.79)	66.67 <sup>cde</sup> (54.74)	63.33 <sup>bcd</sup> (52.73)	56.67 <sup>bcde</sup> (48.83)	50.00 <sup>abcd</sup> (45.00)	46.67 <sup>abc</sup> (43.09)
11	T <sub>11</sub>	90.00 <sup>a</sup> (71.57)	76.67 <sup>bcd</sup> (61.12)	80.00 <sup>ab</sup> (63.44)	73.33 <sup>ab</sup> (58.91)	66.67 <sup>abc</sup> (54.74)	60.00 <sup>abc</sup> (50.77)	53.33 <sup>abc</sup> (46.91)
12	T <sub>12</sub>	90.00 <sup>a</sup> (71.57)	86.67 <sup>a</sup> (68.59)	83.33 <sup>a</sup> (65.90)	76.67 <sup>a</sup> (61.12)	73.33 <sup>a</sup> (58.91)	66.67 <sup>a</sup> (54.74)	60.00 <sup>a</sup> (50.77)
13	T <sub>13</sub>	76.67 <sup>b</sup> (52.73)	60.00 <sup>e</sup> (50.77)	56.67 <sup>e</sup> (48.83)	50.00 <sup>d</sup> (45.00)	43.33 <sup>e</sup> (41.17)	36.63 <sup>d</sup> (37.25)	30.00 <sup>d</sup> (33.21)
14	T <sub>14</sub>	0.00 <sup>c</sup> (0.00)	0.00 <sup>f</sup> (0.00)	0.00 <sup>f</sup> (0.00)	0.00 <sup>e</sup> (0.00)	0.00 <sup>f</sup> (0.00)	0.00 <sup>e</sup> (0.00)	0.00 <sup>e</sup> (0.00)
	General mean	77.38	68.33	64.76	59.76	55.00	49.76	45.24
	F	Sig.	Sig.	Sig.	Sig.	Sig.	Sig.	Sig.
	Sig.	0.000	0.000	0.000	0.000	0.000	0.000	0.000
	SE(m)±	1.85	1.74	1.62	1.65	1.52	1.43	1.36
	C.D.	5.40	5.06	4.73	4.81	4.43	4.16	3.96
	C.V.	5.21	5.47	5.33	5.76	5.63	5.65	5.73

DAS: Days After Storage, Values are the means of three replications,

Values in parentheses are angular transformed values

Means in a column followed by same superscript are not significantly different according to DMRT at P≤ 0.05.

SE(m)± = Standard error of mean C.V. = Coefficient of variation C.D. = Critical Difference at 1% level of significance

T<sub>1</sub>: Mineral oil culture of *B. bassiana* stored at Room temperature (28 ± 3°C), T<sub>2</sub>: Mineral oil culture of *B. bassiana* stored at Refrigerated temperature (4°C), T<sub>3</sub>: Mineral oil culture of *B. bassiana* stored at freezing temperature (-20°C), T<sub>4</sub>: Liquid paraffin oil culture of *B. bassiana* stored at Room temperature (28 ± 3°C), T<sub>5</sub>: Liquid paraffin oil culture of *B. bassiana* stored at Refrigerated temperature (4°C), T<sub>6</sub>: Liquid paraffin oil culture of *B. bassiana* stored at freezing temperature (-20°C), T<sub>7</sub>: Rice bran oil culture of *B. bassiana* stored at Room temperature (28 ± 3°C), T<sub>8</sub>: Rice bran oil culture of *B. bassiana* stored at Refrigerated temperature (4°C), T<sub>9</sub>: Rice bran oil culture of *B. bassiana* stored at freezing temperature (-20°C), T<sub>10</sub>: Sesamum oil culture of *B. bassiana* stored at Room temperature (28 ± 3°C), T<sub>11</sub>: Sesamum oil culture of *B. bassiana* stored at Refrigerated temperature (4°C), T<sub>12</sub>: Sesamum oil culture of *B. bassiana* stored at freezing temperature (-20°C), T<sub>13</sub>: Unformulated culture of *B. bassiana* (Not preserved in oil), T<sub>14</sub>: Untreated control.




Fig. 3. Viability of *Beauveria bassiana* conidia stored under different temperature regimes after formulating as oil cultures (2023-2024)






Fig. 4. Virulence of *Beauveria bassiana* conidia stored under different temperature regimes after formulating as oil cultures against third instar larvae of *Spodoptera litura* under laboratory conditions (2023-2024)

In the investigations after thirty days of storage, the highest mean per cent conidial viability of 82.14% was recorded with Sesamum oil cultures stored at -20°C (T<sub>12</sub>). The treatment T<sub>11</sub> (80.71%) was on par with T<sub>6</sub> (79.28%). The treatments T<sub>9</sub> (76.42%), T<sub>8</sub> (75.71%) and T<sub>3</sub> & T<sub>5</sub> (74.28%) were on par with each other. These were followed by the treatments T<sub>2</sub> and T<sub>4</sub> & T<sub>10</sub> (70.71%). Comparatively lower conidial viability of 67.85% was observed in oil cultures formulated using Mineral oil (T<sub>1</sub>) and Rice Bran oil at room temperature (T<sub>7</sub>). The unformulated culture (T<sub>13</sub>) recorded a conidial viability of 61.43% (Table. 1). The observations of larval mortalities of *Beauveria bassiana* oil cultures after being stored for 30 days revealed that the highest per cent larval mortality of 86.67% was observed in T<sub>12</sub> followed by T<sub>6</sub> (83.33%) and T<sub>9</sub> (80.00%). The next superior treatments T<sub>3</sub>, T<sub>8</sub> & T<sub>11</sub> (76.67%) and T<sub>5</sub> (73.33%) were on par with each other. These were followed by the treatments T<sub>2</sub>, T<sub>4</sub> & T<sub>10</sub> (70.00%) and T<sub>1</sub> & T<sub>7</sub> (66.67%). Comparatively lower mortality of 60.00% was observed in T<sub>13</sub>. The per cent larval mortality in untreated control was 0.00% (Table. 2).

Among the treatments after sixty days of storage, the treatment (T<sub>12</sub>) Sesamum oil cultures stored at -20°C was recorded with the highest mean per cent conidial viability of 79.28%. T<sub>11</sub> (76.42%) was on par with T<sub>9</sub> (75.71%). The next better treatment was T<sub>6</sub> (74.28%). The treatments T<sub>3</sub> & T<sub>5</sub> (70.71%), T<sub>8</sub> (69.28%) and T<sub>4</sub> (67.85%) were on par with each other. These were followed by the treatments T<sub>2</sub> & T<sub>10</sub> (65.71%), T<sub>7</sub> (64.28%) and T<sub>1</sub> (61.43%) were on par with each other. The unformulated culture (T<sub>13</sub>) recorded a conidial viability of 58.57% (Table. 1). After 60 days of storage, highest mean per cent larval mortality of 83.33% was observed in T<sub>12</sub> followed by T<sub>11</sub> (80.00%), T<sub>6</sub> & T<sub>9</sub> (76.67%), T<sub>3</sub>, T<sub>5</sub> & T<sub>8</sub> (70.00%) and T<sub>2</sub>, T<sub>4</sub> & T<sub>10</sub> (66.67%). The treatments T<sub>7</sub> (63.33%) and T<sub>1</sub> (60.00%) were on par with each other. Comparatively lower mortality of 56.67% was observed in T<sub>13</sub>. There was no larval mortality recorded in untreated control T<sub>14</sub> (Table. 2).

It was observed that among all the treatments after ninety days of storage, the treatment T<sub>12</sub> was recorded with the highest mean per cent conidial viability of 75.71% and it was on par with T<sub>11</sub> (74.28%) and T<sub>9</sub> (73.57%). The next superior treatment was T<sub>6</sub> (70.71%). The treatments T<sub>3</sub> & T<sub>5</sub> (65.00%) was on par with T<sub>8</sub> (65.71%). T<sub>10</sub> (63.57%), T<sub>2</sub> (62.87%) and T<sub>4</sub> (61.43%) were on par with each other. These were followed by the

treatments, T<sub>7</sub> & T<sub>1</sub> (57.87%). The unformulated culture (T<sub>13</sub>) recorded a conidial viability of 51.43% (Table. 1). The results of mean per cent larval mortality after ninety days of storage of oil cultures revealed that the highest per cent larval mortality was recorded in T<sub>12</sub> (76.67%), followed by T<sub>9</sub> & T<sub>11</sub> (73.33%). The treatments T<sub>6</sub> (70.00%) and T<sub>3</sub> & T<sub>5</sub> (66.67%) were on par with each other. These treatments were followed by T<sub>8</sub> & T<sub>10</sub> (63.33%) and T<sub>2</sub> & T<sub>4</sub> (60.00%) which were on par with each other. Comparatively lower per cent larval mortalities were observed in T<sub>1</sub> & T<sub>7</sub> (56.67%) and T<sub>13</sub> (50.00%). The per cent larval mortality in untreated control was 0.00% (Table. 2)

The data on observation of mean per cent conidial viability after 120 DAS among all the treatments, T<sub>12</sub> was recorded with the highest mean per cent conidial viability of 74.28%. T<sub>11</sub> (67.85%) was on par with T<sub>9</sub> (69.28%). The next superior treatment T<sub>6</sub> (65.71%), T<sub>3</sub> (64.28%) and T<sub>5</sub> (61.43%) were on par with each other. The treatments T<sub>2</sub> & T<sub>8</sub> (58.57%), T<sub>10</sub> (57.87%) and T<sub>4</sub> & T<sub>7</sub> (54.29%) were statistically indifferent from each other. These were followed by the treatment T<sub>1</sub> (51.43%). The unformulated culture (T<sub>13</sub>) recorded a conidial viability of 45.71% (Table. 1). The observations after 120 days of storage of *B. bassiana* oil cultures revealed that maximum of 73.33% mean per cent larval mortality was recorded in T<sub>12</sub> followed by T<sub>9</sub> (70.00%) and T<sub>6</sub> & T<sub>11</sub> (66.67%). T<sub>3</sub> (63.33%) was on par with T<sub>5</sub> (60.00%). These treatments were followed by T<sub>2</sub>, T<sub>8</sub> and T<sub>10</sub> (56.67%), T<sub>4</sub> & T<sub>7</sub> (53.33%) followed by T<sub>1</sub> (50.00%) and T<sub>13</sub> (43.33%). There was no mortality recorded in untreated control T<sub>14</sub> (Table. 2).

The results after 150 days of storage of different oil cultures of *Beauveria bassiana* revealed that among the treatments T<sub>12</sub> was recorded with the highest mean per cent conidial viability of 67.85%, followed by T<sub>9</sub> (64.28%). T<sub>6</sub> (63.75%) was on par with T<sub>11</sub> (62.87%). The treatment T<sub>3</sub> & T<sub>5</sub> (58.57%) on par with T<sub>8</sub> (55.71%). The treatments T<sub>2</sub> (51.43%) and T<sub>10</sub> & T<sub>4</sub> (49.29%) were statistically indifferent. These were followed by the treatments T<sub>7</sub> (48.57%) and T<sub>1</sub> (44.29%). The unformulated culture (T<sub>13</sub>) recorded a conidial viability of 39.29% (Table. 1). Storage of oil cultures for 150 days in the laboratory showed that highest mean per cent larval mortality of 66.67% was recorded in T<sub>12</sub> followed by T<sub>6</sub> & T<sub>9</sub> (63.33%). T<sub>11</sub> (60.00%) and T<sub>3</sub> & T<sub>5</sub> (56.67%) were on par with each other. The treatments T<sub>8</sub> (53.33%) and T<sub>2</sub>, T<sub>4</sub> & T<sub>10</sub> (50.00%) were

significantly indifferent with each other. These were followed by T<sub>7</sub> (46.67%), T<sub>1</sub> (43.33%) and T<sub>13</sub> (36.63%). The per cent larval mortality in untreated control was 0.00%. (Table. 2).

The results of the percent viability after 180 days of storage clearly shows that, the highest mean per cent conidial viability of 62.87% was recorded with T<sub>12</sub> followed by T<sub>6</sub> (61.43%), T<sub>9</sub> (58.57%), T<sub>3</sub> & T<sub>11</sub> (54.29%) and T<sub>5</sub> & T<sub>8</sub> (51.43%). The treatments T<sub>2</sub> (48.57%), T<sub>10</sub> (47.85%) and T<sub>4</sub> (45.71%) were on par with each other. These were followed by the treatments T<sub>7</sub> (44.29%) and T<sub>1</sub>(41.43%). The unformulated culture (T<sub>13</sub>) recorded a conidial viability of 32.85% (Table. 1). The data after 180 days of storage of *Beauveria bassiana* oil cultures clearly revealed that highest mean larval mortality of 60.00% was recorded with T<sub>12</sub> & T<sub>6</sub> followed by T<sub>9</sub> (56.67%). The treatments T<sub>3</sub> & T<sub>11</sub> (53.33%), T<sub>5</sub> & T<sub>8</sub> (50.00%) and T<sub>2</sub> & T<sub>10</sub> (46.67%) were on par with each other. These were followed by the treatments T<sub>4</sub> & T<sub>7</sub> (43.33%) followed by T<sub>1</sub> (40.00%) and T<sub>13</sub> (30.00%). There was no larval mortality recorded in untreated control T<sub>14</sub> (Table. 2).

Comparatively higher mean per cent conidial viabilities of *Beauveria bassiana* oil cultures were observed in the treatments T<sub>12</sub>, T<sub>6</sub>, T<sub>9</sub> and T<sub>11</sub>, from the day of preparation to 180 days after storage. The viability varied as follows: T<sub>12</sub> (87.85% to 62.87%), T<sub>6</sub> (83.57 to 61.43%), T<sub>9</sub> (80.71 to 58.57%) and T<sub>11</sub> (87.85 to 54.29%). Comparatively lower per cent conidial viability were recorded in T<sub>1</sub> (77.41 to 41.43%) and T<sub>13</sub> (74.28 to 32.85%). The highest mean per cent larval mortalities were observed in Treatments T<sub>12</sub>, T<sub>6</sub>, T<sub>9</sub> and T<sub>11</sub> when treated against third instar *S. litura* larvae at monthly intervals. The mean per cent larval mortalities ranged as follows: T<sub>12</sub> (90.00% to 60.00%), T<sub>6</sub> (86.67 to 60.00%), T<sub>9</sub> (83.33 to 56.67%) and T<sub>11</sub> (90.00 to 53.33%). Comparatively lower per cent larval mortalities were recorded in T<sub>1</sub> (80.00 to 40.00%) and T<sub>13</sub> (76.67 to 30.00%).

The pathogenicity levels correlated with conidial viabilities, showing a gradual decrease in both viability and virulence over the 180-days storage period.

From the present study, it can be concluded that the performance of *Beauveria bassiana* was best with oil formulations compared to unformulated crude cultures. Among the oils used, Sesamum oil formulations were the most effective followed

by Liquid paraffin oil, Rice bran oil and mineral oil. Considering the temperature conditions for storage, the formulations stored at freezing temperature (-20°C) performed better to those stored at Refrigerated temperature (4°C) and Room temperature (28 ± 3°C).

Sesamum oil cultures had been found out to be best among all oil cultures, this may be due to Sesquiterpene present in it which have insecticidal properties that serve as a synergist for pyrethroid insecticides [30-32].

The performance of oil cultures was best at lower temperatures compared to room temperature, which may be due to reduced metabolism and growth at lower temperatures thereby maintaining viability for an extended period in such lower temperatures. Murugasridevi *et al.* [26] prepared oil-based formulation by dissolving 1 g of pure conidia (10<sup>10</sup> conidia g<sup>-1</sup>) of *B. bassiana* (Bb 112) in 100 ml of paraffin oil, along with adjuvants like polyethylene glycol and tween 80 to enhance the efficacy of the formulation and stored it under different storage conditions *viz.* ambient temperature (28 ± 3°C) and refrigerated condition (4°C). Shelf-life assessment showed that the formulation stored under ambient temperature and refrigerated condition recorded 587.00 and 590.77×10<sup>8</sup> CFU ml<sup>-1</sup>, respectively on the day of preparation and 167.66 and 216.66×10<sup>8</sup> CFU ml<sup>-1</sup>, respectively after 24 weeks.

The results were in agreement with Lakshmidevi *et al.* [33] prepared 13 types of plant oil-based bio-formulation of *Beauveria bassiana* (TBb8 strain) using corn, soyabean, castor, rice bran, mustard, mahuva, pinnai, neem, groundnut, palm, coconut and gingelly oils with different combinations and tested its efficacy against the lepidopteran fruit borer *Helicoverpa armigera* (Hubner). Among 13 oils, corn oil was found superior by maintaining a population of 7.0×10<sup>9</sup> cfu/ml after 210 days of storage. Oil formulation of TBb8 isolate (1×10<sup>8</sup> spore/ml) treated *H. armigera* showed larval mortality of 70.0%, pupation period of 13.7 days, pupal malformation of 17.9% and adult malformation was recorded as 16.2%.

Sandhu *et al.* [27] evaluated the effects of temperature on conidial viability and virulence of *B. bassiana* against third instar *Helicoverpa armigera* (Hubner) larvae over a 24-month period. He stored unformulated conidia of *B. bassiana* at five different temperatures (0°C,

10°C, 20°C, 30°C and 40°C). Based on the experimental results they reported that conidia survived longest at lower temperatures (0–20°C). At higher temperatures, (30–40°C) conidia did not survive.

Oliveira *et al.* [34] evaluated the effect of preservation of *B. bassiana* isolates for one year on growth, production and viability of spores as well as macro- and micro-morphology by using 3 storage methods *viz.* freezing at -20°C in an aqueous glycerol solution; freeze-drying (lyophilized) and maintaining at ambient temperature with continual sub-culturing in PDA medium. They reported that glycerol-freeze method at -20°C is the most suitable method for long-term preservation of fungal species.

#### 4. CONCLUSIONS

The results indicated that *Beauveria bassiana* could be a potent biocontrol agent for managing lepidopteran pests. The combinations of cost-effective compatible oils with entomopathogenic fungal agents and storage under controlled temperature conditions presents a promising strategy in enhancing the effectiveness of fungal agents compared to unformulated crude culture. This approach not only enhances the practical application but also supports sustainable pest management practices by providing a reliable and environmentally friendly alternative to chemical pesticides. Further research should be conducted in this line at field level to identify the most effective oil type and concentrations suiting different environment conditions and pest targets, as well as optimizing the application methods to ensure uniform coverage and maximum reach to pests. Additionally, research on compatibility between entomopathogenic fungal oil cultures and other biocontrol agents or agricultural practices will improve its effectiveness and reliability.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### ACKNOWLEDGEMENTS

The authors are sincerely thankful to the Department of Entomology, S.V. Agricultural

college, Tirupati, ANGRAU, India for providing infrastructure and financial support.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. FAO. Pesticides Use, Pesticides Trade and Pesticides Indicators-Global, Regional and Country Trends, 1990–2020. FAOSTAT Analytical Briefs. 2022; 46.
2. Dong H, Zhu KH, Zhao Q, Bai XP, Zhou JC, Zhang LS. Morphological defense of the egg mass of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) affects parasitic capacity and alters behaviours of egg parasitoid wasps. Journal of Asia-Pacific Entomology. 2021;24(3):671-678.
3. Waqas MS, Shi Z, Yi TC, Xiao R, Shoaib AA, Elabasy AS, Jin DC. Biology, ecology, and management of cotton mealybug *Phenacoccus solenopsis* Tinsley (Hemiptera: Pseudococcidae). Pest Management Science. 2021;77(12):5321-5333.
4. Guo J, Wu S, Zhang F, Huang C, He K, Babendreier D, Wang Z. Prospects for microbial control of the fall armyworm *Spodoptera frugiperda*: a review. Bio Control. 2020; 65:647-662.
5. Yan-li Z, Hui D, Li-sheng Z, Zu-min G, Jin-cheng Z. High virulence of a naturally occurring entomopathogenic fungal isolate, *Metarhizium (Nomuraea) rileyi*, against *Spodoptera frugiperda*. Journal of Applied Entomology. 2022;146(6):659-665.
6. Ummidi VRS, Vadlamani P. Preparation and use of oil formulations of *Beauveria bassiana* and *Metarhizium anisopliae* against *Spodoptera litura* larvae. African Journal of Microbiology Research. 2014;8 (15):1638-1644.
7. Sharma A, Sharma S, Yadav PK. Entomopathogenic fungi and their relevance in sustainable agriculture: A review. Cogent Food & Agriculture. 2023; 9(1).
8. Bateman RP, Douro-Kpindou OK, Kooyman C, Lomer C, Ouambama Z. Some observations on the dose transfer of mycoinsecticide sprays to desert locusts. Crop Protection. 1998;17(2):151-158.

9. Ramanujam B, Rangeshwaran R, Sivakumar G, Mohan M, Yandigeri MS. Management of insect pests by microorganisms. Proceedings of the Indian National Science Academy. 2014;80(2): 455-471.
10. Gebremariam A, Chekol Y, Assefa F. Extracellular enzyme activity of entomopathogenic fungi, *Beauveria bassiana* and *Metarhizium anisopliae* and their pathogenicity potential as a bio-control agent against whitefly pests, *Bemisia tabaci* and *Trialeurodes vaporariorum* (Hemiptera: Aleyrodidae). BMC Research Notes. 2022;15(1):117.
11. Seema Y, Neeraj T, Krishan K. Mass production of entomopathogens *Beauveria bassiana* and *Metarhizium anisopliae* using rice as a substrate by diphasic liquid-solid fermentation technique. International Journal of Advanced Biological Research. 2013;3(3):331-335.
12. Beys-da-Silva WO, Santi L, Berger M, Calzolari D, Passos DO, Guimarães JA, Moresco JJ, Yates JR. Secretome of the biocontrol agent *Metarhizium anisopliae* induced by the cuticle of the cotton pest *Dysdercus peruvianus* reveals new insights into infection. Journal of Proteome Research. 2014;13(5):2282-2296.
13. de Faria MR, Wraight SP. Mycoinsecticides and mycoacaricides: a comprehensive list with worldwide coverage and international classification of formulation types. Biological control. 2007; 43(3):237-256.
14. Money NP. Fungi and biotechnology. The fungi. 2016;401-424.
15. Singh HB, Keswani C, Ray S, Yadav SK, Singh SP, Singh S, Sarma BK. *Beauveria bassiana*: biocontrol beyond lepidopteran pests. Biocontrol of Lepidopteran Pests: Use of Soil Microbes and their Metabolites. 2015;219-235.
16. Wang Q, Xu L. Beauvericin, a bioactive compound produced by fungi: a short review. Molecules. 2012;17(3):2367-2377.
17. Rachappa V, Lingppa S, Patil RK, Hugar PS. Suitability of media and containers for mass production of *Metarhizium anisopliae*. Karnataka Journal of Agricultural Sciences. 2005;18(3): 680.
18. Inyang EN, McCartney HA, Oyejola B, Ibrahim L, Archer SA. Effect of formulation, application and rain on the persistence of the entomogenous fungus *Metarhizium anisopliae* on oilseed rape. Mycological Research. 2000;104(6):653-661.
19. Devi PV, Prasad YG. Compatibility of oils and antifeedants of plant origin with the entomopathogenic fungus *Nomuraea rileyi*. Journal of invertebrate pathology. 1996;68(1):91-93.
20. Bateman RP, Carey M, Moore DE, Prior C. The enhanced infectivity of *Metarhizium flavoviride* in oil formulations to desert locusts at low humidities. Annals of Applied Biology. 1993;122(1): 145-152.
21. Batta YA. Production and testing of novel formulations of the entomopathogenic fungus *Metarhizium anisopliae* (Metschnikoff) Sorokin (Deuteromycotina: Hyphomycetes). Crop Protection. 2003;22(2):415-422.
22. Moore D, Bridge PD, Higgins PM, Bateman RP, Prior C. Ultra-violet radiation damage to *Metarhizium flavoviride* conidia and the protection given by vegetable and mineral oils and chemical sunscreens. Annals of Applied Biology. 1993;122(3): 605-616.
23. Humber RA. Fungi: preservation of cultures. In Manual of techniques in insect pathology. Academic Press. 1997;269-279.
24. López Lastra CC, Hajek AE, Humber RA. Comparing methods of preservation for cultures of entomopathogenic fungi. Canadian Journal of Botany. 2002;80 (10): 1126-1130.
25. Humber RA. Preservation of entomopathogenic fungal cultures. Manual of techniques in invertebrate pathology. 2012;2:317-328.
26. Murugasidevi K, Jeyarani S, Kumar SM. Evaluation of shelf life and pathogenicity of oil-based formulation of *Beauveria bassiana* against chilli thrips and mites. Journal of Entomological Research. 2021;45(4):679-685.
27. Sandhu SS, Rajak RC, Agarwal GP. Studies on prolonged storage of *Beauveria bassiana* conidia: effects of temperature and relative humidity on conidial viability and virulence against chickpea borer, *Helicoverpa armigera*. Biocontrol Science and Technology. 1993;3(1):47-53.
28. Saheb YP, Manjula KK, Devaki Devi RSJ, Reddy BR, Anokhe A. Evaluation of *Metarhizium (Nomuraea) rileyi* Rice Bran Oil Formulations against 3<sup>rd</sup> instar *Spodoptera litura* under Laboratory. Biological Forum – An International Journal. 2021;13(3):601-607.

29. Torrado-León E, Montoya-Lerma J, Valencia-Pizo E. Sublethal effects of *Beauveria bassiana* (Balsamo) Vuillemin (Deuteromycotina: Hyphomycetes) on the whitefly *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) under laboratory conditions. *Mycopathologia*. 2006;162: 411-419.
30. Wei P, Zhao F, Wang Z, Wang Q, Chai X, Hou G, Meng Q. Sesame (*Sesamum indicum* L.): A comprehensive review of nutritional value, phytochemical composition, health benefits, development of food, and industrial applications. *Nutrients*. 2022;14(19):4079.
31. Baker BP, Grant JA. Active ingredients eligible for minimum risk pesticide use: Overview of the profiles; 2018
32. Sirato-Yasumoto S, Katsuta M, Okuyama Y, Takahashi Y, Ide T. Effect of sesame seeds rich in sesamin and sesamol on fatty acid oxidation in rat liver. *Journal of Agricultural and Food Chemistry*. 2001; 49(5):2647-2651.
33. Lakshmidēvi P, Gopalakrishnan C, Antony RS. Bio-formulation based on plant oil of the future *Beauveria bassiana* (Balsamo) for fruit borer control in tomatoes. *Journal of Entomology and Zoology Studies*. 2020;8:200-206.
34. Oliveira I, Pereira JA, Bento A, Baptista P. Viability of *Beauveria bassiana* isolates after storage under several preservation methods. *Annals of Microbiology*. 2011;61 (2):339-344.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/120760>