



Character Association and Path Coefficient Analysis in Upland Rice (*Oryza sativa* L.) for Grain Yield and Quality Characters

Jalla Rajasekhar ^{a*}, G. R. Lavanya ^{a#}, Gattikoppula Purushotham ^{a‡}
and D. Praveen Kumar ^{a‡}

^a Department of Genetics and Plant Breeding, Faculty of Agriculture, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj- 211007, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i2331605

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92570>

Original Research Article

Received 02 August 2022
Accepted 04 October 2022
Published 06 October 2022

ABSTRACT

The present investigation was undertaken to evaluate Twenty Five Upland Rice genotypes. There are various aspects to move up rice yield, such as creating of irrigation work, development of rice ecosystems, soil conditions, traditional practices, and breeding of high yielding genotypes. Among them, it seems that the most effective and economic way available is to develop different rice ecosystems especially Upland Rice Genotypes. High to moderate phenotypic and genotypic coefficient of variation were observed for number of panicles per hill and number of tillers per hill. High heritability coupled with high genetic advance as percent of mean was observed for number of panicles per hill, number of tillers per hill, grain yield per hill, test weight and flag leaf length, indicating that these traits are most probably under the control of additive gene action and hence these traits can be fixed by proper selection. Days to 50% flowering, plant height, number of panicles per hill, flag leaf width, spikelets per panicle, days to maturity, biological yield per hill, harvest index and test weight, exhibited positive significant correlation with grain yield per hill at both genotypic and phenotypic levels, indicating selection based on these traits could be more

[‡] M. Sc Scholar;

[#] Associate Professor;

*Corresponding author: E-mail: jallarajasekhar3@gmail.com;

effective to maximize grain yield. Days to 50% flowering, biological yield per hill and harvest index had positive direct effect on grain yield per hill. Therefore, selection program should be based on these traits for crop improvement with high yielding genotypes in Upland rice.

Keywords: *Upland rice; genetic variability; heritability; genetic advance; correlation coefficient analysis; path coefficient analysis.*

1. INTRODUCTION

"Drought is one of the most known limiting factors of crop cultivation leading to unstable food security. Osmolyte like proline helps the plants to escape from drought stress by protecting from dehydration rapidly. Deep rooted plants show greater drought avoidance capacity than shallow-rooted plants. It has also been reported that drought resistant upland rice varieties remained tall and generally have low tillering capacity" [1]. "Rice cultivation consumes about 30% of all fresh water used worldwide. In Asia, flood irrigated rice consumes more than 45% of total fresh water used" [2]. "By 2025, 15 out of 75 millions hectare of Asia's flood irrigated rice crop will experience water shortage" [3]. "With increasing water scarcity, the sustainability, food production, and ecosystem services of rice fields are threatened" [4]. To overcome the water scarcity problem upland is an important system of rice culture and constitutes about one sixth of the world's total rice land. Direct seeding rice, a common practice before green revolution in India, is once again became popular because of its potential to save water and labour [5]. "Upland rice is also called autumn rice. The study of genetic variability in any crop would help in the genetic improvement of yield and desirable characters. It will facilitate the identification of proper genotypes for a particular agro-climate zone. Heritability along with the genetic advance would be more useful tool in predicting the resultant effect from selection of the best genotypes for yield and some of its components in rice" [6]. "Correlation and genetic variability analysis leads us to a clear understanding of the genetic association of various plant traits and their contribution to yield. correlation coefficients are generally employed to determine the relation of grain yield and yield components" [7]. "Path coefficient analysis is an efficient statistical technique specially designed to quantify the interrelationship of different components and their direct and indirect effects on grain yield" [8]. "The information on relative direct and indirect contribution of each component character toward yield will help breeders to formulate the effective criteria in selecting desirable genotypes in early

segregating populations" [9]. In view of this, the present study was planned to determine the genetic variability, heritability, genetic advance, correlation coefficients and direct and indirect contribution of twenty five upland rice genotypes yield and yield attributing traits on grain yield.

1.1 Objectives

1. To estimate genetic variability, heritability and genetic advance for yield and its related traits in rice cultivars.
2. To study the nature of character association among yield and grain yield attributing traits.
3. To assess the direct and indirect contribution of yield attributing traits on grain yield.

2. MATERIALS AND METHODS

Twenty five upland rice genotypes (Table 1) collected from SHUATS were sown in the randomized block design with three replications at the Crop Research Farm of the Department of Genetics and Plant Breeding, Sam Higginbottom University of Agriculture technology and Sciences, Prayagraj, U.P during Kharif - 2021. In each replication, two to three seedling were sowed per hill with 20x10 cm spacing. The observations were recorded on five randomly selected plants from each plot for days to 50 percent flowering, plant height, number of tillers per hill, number of panicles per hill, panicle length, number of spikelets per panicle, spikelet fertility, flag leaf length, flag leaf width, test weight, days to maturity, biological yield per hill, harvest index and grain yield per hill. As per established methods, data were analysed to determine Analysis of variance [10]. The statistical and biometrical analysis of GCV and PCV were calculated by formula given by Burton (1952), [11]. Heritability in broad sense (h^2) by Burton and de Vane (1953), [12]. The expected genetic gain was calculated using the procedure given by Johnson et al.(1955), [13]. The correlation coefficient and path coefficient were worked out as for the method recommended by Al-Jibouri et al.(1958) and Deway and Lu (1959), [14] respectively, the estimated values were

compared with table values of the correlation coefficient to test the significance of the correlation coefficient prescribed by Fisher and Yates (1963), [15].

2.1 Layout Description

Season: *Kharif* - 2021

Crop: Upland Rice (*Oryza sativa* L.)

Number of genotypes : 24+1 check

Experimental design : Randomized Block Design

Gross experimental area : 240 sq.m

Net experimental area : 210 sq.m

Row to row distance :20 cm

Plant to plant spacing :10cm

Date of sowing: 3-8-2021

Fertilizer dose: N:P:K @ 120:60:50 kg/ha

Date of Harvesting: 15-11-2021

Recommended package of practice were followed to raise a healthy crop

3. RESULTS AND DISCUSSION

3.1 Analysis of Variance

Analysis of Variance revealed existence of sufficient genetic variability among all the genotypes (Table 3). The mean sum of squares for fifteen quantitative traits of 25 upland rice genotypes reported high significant differences among the genotypes for all the traits at 1% level of significance. This indicates that there was an ample scope for selection of promising lines for yield and its components. The presence of large amount of variability might be due to diverse source of materials as well as environmental influence affecting the phenotypes. Similar results have been reported by Girma et al. [16] and Magar et al. [17].

3.1.1 Genotypic and Phenotypic variance

Wide range of Genotypic and Phenotypic coefficient of variance were observed in the experimental material for all the traits (Table 4). The higher phenotypic coefficient of variation was recorded for number of panicles per hill (24.10) followed by number of tillers per hill (22.28). And higher genotypic coefficient of variation was observed for number of panicles per hill (22.92) followed by number of tillers per hill (21.44). it is evident that phenotypic coefficient of variation values were higher than genotypic coefficient of variation for all the traits under study indicating the influence of environment on studied characters. The results are parallel with the findings of Kumar et al. [18], Khatun et al. [19], Chandra et al. [20], Anjaneyulu et al. [21].

3.1.2 Heritability

“Estimation of genotypic coefficient of variation reflect the total amount of genotypic variability, the proportion of this genotypic variability, which transmitted from parents to the progeny, is reflected through heritability”. Lush (1949), [22], gave “the concept of broad sense heritability, which is the ratio of genotypic variance (VG) to phenotypic variance (VP). It determines efficiency with which we can utilize the genotypic variability in breeding programs”. Burton (1952) suggested that “genetic variation along with heritability will give better idea about expected efficiency of selection. In the present study estimates of broad sense heritability are computed which includes both additive and non-additive gene effects. Higher values of broad sense heritability recorded for the traits (Table 4), Number of tillers per hill (92.62) followed by number of panicles per hill (90.43), test weight (84.67), grain yield per hill (84.03), flag leaf length (83.34), spikelets per panicle (81.88), plant height (61.89) these characters are less influenced by environmental effects and selection on the basis of phenotypic performance of genotypes would be more efficient in further improvement of these traits”. Similar results have been reported by Tiwari et al. [23] and Pratap et al. [24].

3.1.3 Mean performance of superior genotypes for quality parameters based on grain yield

According to the above (Table 2), Among 25 accessions SHUATS UPR –33 had highest kernel length before cooking (6.37), L/B ratio (3.76) and kernel length after cooking (7.90). SHUATS UPR – 31 had highest mean values for kernel width after cooking (2.97) and elongation ratio (1.25). SHUATS UPR – 34 exhibited highest values for alkali spreading value (6.52) and hulling percentage (73.4).SHUATS UPR-43 (25.04) was found to be highest in amylose content and SHUATS UPR-30 (76.00) was found to be highest in gel consistency.

3.1.4 Genetic Advance

“In present study (Table 4), Estimates of genetic advance ranged from (0.08) flag leaf width to (32.48) spikelet per panicle. High genetic advance was recorded for spikelet per panicle (32.48).). Moderate genetic advance was recorded for plant height (15.40). Traits such as flag leaf length (8.69), test weight (5.94), harvest index (4.73), biological yield (3.97), spikelet

fertility (3.45), grain yield per hill (3.24), days to maturity (2.94), days to 50% flowering (2.91), panicle length (1.64), number of tillers per hill (2.70), number of panicles per hill (2.44) and flag leaf width (0.08) exhibited low genetic advance. High to Moderate estimates of genetic advance for the traits, Spikelets per panicle and plant

height suggested that these parameters were under the control of additive genetic effects and selection at early segregating generation for these traits will prove beneficial for selecting superior varieties of upland rice". Similar results were reported by Longjam and Singh [25] and Singh et al. [26].

Table 1. List of Experimental Materials

Sr. No.	Advanced breeding lines	Cross combination	Grain type
1.	SHUATS UPR-25	IR82589-B-B-121-3 X NDR-97 (P2)	Medium bold
2.	SHUATS UPR-26	IR82589-B-B-121-3 X NDR-97 (P3)	Medium bold
3.	SHUATS UPR-27	IR82589-B-B-121-3 X NDR-97 (P5)	Medium bold
4.	SHUATS UPR-28	IR82589-B-B-121-3 X NDR-97 (P7)	Medium bold
5.	SHUATS UPR-29	IR82589-B-B-121-3 X Sahbhagidhan	Medium slender
6.	SHUATS UPR-30	IR82589-B-B-121-3 X Sahbhagidhan	Medium slender
7.	SHUATS UPR-31	IR82589-B-B-121-3 X Sahbhagidhan	Medium slender
8.	SHUATS UPR-32	IR82589-B-B-121-3 X Sahbhagidhan	Medium slender
9.	SHUATS UPR-33	IR82589-B-B-121-3 X Sahbhagidhan	Medium slender
10.	SHUATS UPR-34	IR82589-B-B-121-3 X Sahbhagidhan	Medium slender
11.	SHUATS UPR-35	IR82589-B-B-121-3 X Sahbhagidhan	Long slender
12.	SHUATS UPR-36	IR82639-B-B-200-4 X Sahbhagidhan	Long slender
13.	SHUATS UPR-37	IR82639-B-B-200-4 X Sahbhagidhan	Long slender
14.	SHUATS UPR-38	IR82639-B-B-200-4 X Sahbhagidhan	Long slender
15.	SHUATS UPR-39	IR82639-B-B-200-4 X Sahbhagidhan	Long slender
16.	SHUATS UPR-40	IR82639-B-B-200-4 X Sahbhagidhan	Long slender
17.	SHUATS UPR-41	IR82639-B-B-200-4 X Sahbhagidhan	Long slender
18.	SHUATS UPR-42	IR82639-B-B-200-4 X Sahbhagidhan	Long slender
19.	SHUATS UPR-43	IR83750-B-B-131-1 X NDR-97 (P1)	Long slender
20.	SHUATS UPR-44	IR83750-B-B-131-1 X NDR-97 (P3)	Long slender
21.	SHUATS UPR-45	IR83750-B-B-131-1 X NDR-97 (P4)	Long slender
22.	SHUATS UPR-46	IR83750-B-B-131-1 X NDR-97 (P5)	Medium bold
23.	SHUATS UPR-47	IR83750-B-B-131-1 X NDR-97 (P7)	Medium bold
24.	SHUATS UPR-48	IR83750-B-B-131-1 X Sahbhagidhan	Medium slender
25.	SAHBHAGIDHAN (CHECK-1)		Medium slender

Table 2. Mean performance of superior five genotypes for quality parameters

S.No.	Characters	SHUATS UPR - 34	SHUATS UPR - 33	SHUATS UPR - 43	SHUATS UPR - 31	SHUATS UPR - 30
1.	Kernel length Before Cooking	6.24	6.37	5.68	5.65	6.11
2.	Kernel width Before Cooking	1.76	1.69	2.26	1.75	1.99
3.	Kernel length After Cooking	7.40	7.90	6.86	7.09	7.12
4.	Kernel width After Cooking	2.36	2.78	2.86	2.97	2.81
5.	L/B Ratio	3.54	3.76	1.98	3.22	3.07
6.	Elongation Ratio	1.18	1.24	1.20	1.25	1.16
7.	Gel Consistency	54.33	72.66	65.00	70.66	76.00
8.	Amylose content	21.77	22.61	25.04	24.89	23.82
9.	Alkali spreading value	6.52	5.25	6.15	5.81	2.01
10.	Hulling(%)	73.4	68.5	71.3	65.8	70.2

Table 3. Analysis of Variance for fifteen quantitative traits of 25 Upland rice genotypes

Characters	Mean sum of squares		
	Replication (df=02)	Treatments (df= 24)	Error (df=48)
Days to 50% flowering	28.47	33.43**	14.19
Plant height	28.98	326.79**	55.64
No. of tillers per hill	0.40	5.73**	0.14
No. of Panicles per hill	0.13	4.84**	0.16
Panicle length	4.94	6.25**	1.87
Flag leaf length	8.71	68.45**	4.27
Flag leaf width	0.01	0.01**	0.01
Spikelets per panicle	96.27	977.92**	67.16
Spikelet fertility	47.41	52.88**	23.74
Days to maturity	21.07	27.89**	10.57
Biological yield per hill	1.29	23.6**	4.49
Harvest index	2.33	64.84**	22.94
Test weight	2.70	31.32**	1.78
Grain yield per hill	0.18	9.42**	0.56
Grain yield per ha	0.01	1.02**	0.06

** Significant 1% level of significance

Table 4. Genetic Variability parameters for 15 quantitative characters

Characteristics	GCV	PCV	h ² (bs)	GA	GAM
Days to 50% flowering	3.451	6.185	31.126	2.911	3.966
Plant height	9.122	11.595	61.897	15.408	14.784
No. of tillers per hill	21.448	22.285	92.625	2.706	42.522
No. of Panicles per hill	22.925	24.108	90.43	2.445	44.91
Panicle length	5.284	7.982	43.826	1.648	7.206
Flag leaf length	13.51	14.798	83.348	8.698	25.408
Flag leaf width	7.555	11.28	44.856	0.086	10.423
Spikelets per panicle	13.304	14.702	81.885	32.48	24.8
Spikelet fertility	3.48	6.459	29.031	3.459	3.863
Days to maturity	2.365	3.98	35.309	2.941	2.895
Biological yield per hill	10.567	13.805	58.593	3.979	16.663
Harvest index	8.074	13.125	37.843	4.736	10.232
Test weight	14.025	15.242	84.665	5.948	26.584
Grain yield per hill	15.596	17.013	84.037	3.246	29.452
Grain yield per ha	15.618	17.035	84.052	1.071	29.496

GCV: Genotypic Coefficient of Variation, PCV: Phenotypic Coefficient of Variation, h²(bs): Heritability (broad sense), GA: Genetic Advance, GAM: Genetic Advance as Percent of mean

3.1.5 Genetic advance as % of mean

"The estimation of genetic advance as percent mean helps to understand the type of gene action involved in the expression of a trait. It is classified as low (less than 10%), moderate (10-20%) and high (more than 20%) as proposed by Johnson *et al.* [27]. In present study, the estimates of genetic advance as % of mean varied from (2.89) days to maturity to (44.91) number of panicles per hill. High GAM recorded for number of panicles per hill(44.91)

followed by number of tillers per hill (42.52), grain yield per hill (29.45), test weight (26.58), flag leaf length (25.40) and spikelets per panicle (24.8). High genetic advance as percent of mean coupled with high heritability was recorded for number of tillers per hill followed by number of panicles per hill, flag leaf length, test weight and grain yield per hill indicating that these traits are most probably under the control of additive gene action and hence these traits can be fixed by proper selection". Similar results have been reported by Rashmi *et al.* [28].

3.2 Correlation Coefficient Analysis

Correlation study indicates the degree of interdependence of plant characters, which is an important toll in selection of a pertinent genotype. Therefore, information on association between characters is quite useful to plant breeders to formulate their breeding and selection strategies. At present study, Grain yield per plant showed the positive significant phenotypic association with (Table 5) days to 50% flowering(0.311*), plant height (0.337*), number of tillers per plant (0.312*), Number of panicles per hill (0.472**), flag leaf width (0.426**), spikelets per panicle (0.346*), days to maturity(0.305*), biological yield per hill (0.659**), harvest index(0.638**) and test weight(0.274*) while, it exhibited positive non-significant association with panicle length (0.1589) and flag leaf length(0.2035). Grain yield per hill showed positive significant genotypic association with days to 50% flowering (0.392**), plant height (0.501**), number of tillers per plant (0.333*), number of panicles per hill (0.505**), panicle length (0.294*), flag leaf length (0.311*), flag leaf width (0.649**), spikelets per panicle (0.459**), days to maturity (0.412**), biological yield per hill (0.879**), harvest index (0.849**) and Test weight (0.343*). Thus, days to 50% flowering, plant height, Number of panicles per hill, flag leaf width, spikelets per panicle, days to maturity, biological yield per hill, harvest index, test weight were found to possess positive significant association with grain yield per hill at both genotypic and phenotypic level indicating

selection based on these traits could be more effective to maximize grain yield. similar with the findings of Singh et al. [29] and Srijan et al. [30].

3.3 Path Coefficient Analysis

“Correlation gives only the relation between two variables whereas path coefficient analysis allows separation of the direct effect and their indirect effects through other attributes by partitioning the correlation” (Wright, 1921), [31]. “In the present investigation, the path coefficient analysis at phenotypic level revealed that the characters such as (Table 6) days to 50% flowering (0.0448), plant height(0.0119), number of tillers per hill (0.0176), number of panicles per hill (0.0080), flag leaf width (0.0030), biological yield per hill (0.7419) and harvest index (0.7292) had positive direct effect on grain yield per hill. At the Genotypic level path coefficient analysis based on yield as dependent variable showed that days to 50% flowering (0.0562), panicle length (0.1732), flag leaf length (0.0714), spikelet fertility (0.0612), days to maturity (0.0711), biological yield per hill (0.4685), harvest index(0.5446) revealed positive direct effect on grain yield per hill. Therefore, selection program should be based on these traits for crop improvement with high yielding genotypes in Upland rice. These findings of path Coefficient analysis in upland rice were found to be similar with the findings” of Sharma et al. [32], Mishu et al. [33] and Belete et al. [34].




Fig. 1. Genotypic path coefficient




Fig. 2. Phenotypic path coefficient

Table 5. Phenotypic and Genotypic correlation coefficient for grain yield and its component characters in 25 Upland rice genotypes

		PH	NT/H	NP/H	PL	FLL	FLW	S/P	SF	DM	BY/H	HI	TW	GY/H
DF50	P	-0.0330	-0.1065	-0.0369	-0.288*	-0.0065	0.1936	0.379**	-0.234*	0.508**	0.1979	0.1633	-0.0217	0.311*
	G	-0.0299	-0.247*	-0.0817	-0.445**	0.0518	0.756**	0.902**	-0.650**	0.824**	0.1759	0.536**	-0.0722	0.392**
PH	P	1	0.0471	0.1752	0.328*	0.431**	0.1387	0.1843	-0.2082	0.0683	0.298*	0.1390	-0.0227	0.337*
	G	1	0.0663	0.230*	0.618**	0.621**	0.364*	0.307*	-0.537**	0.1765	0.724**	0.0709	-0.0410	0.501**
NT/H	P		1	0.845**	0.413**	0.0964	0.423**	-0.334*	-0.1281	-0.0531	0.1928	0.1944	0.0528	0.312*
	G		1	0.904**	0.648**	0.0856	0.634**	-0.354*	-0.1938	-0.1825	0.291*	0.249*	0.0477	0.333*
NP/H	P			1	0.446**	0.1310	0.532**	-0.256*	-0.1249	-0.0095	0.340*	0.257*	0.1206	0.472**
	G			1	0.755**	0.1404	0.807**	-0.290*	-0.249*	0.0441	0.529**	0.292*	0.1538	0.505**
PL	P				1	0.1848	0.303*	-0.1537	0.0147	-0.1788	0.2112	0.0030	0.0785	0.1589
	G				1	0.338*	0.592**	-0.1160	0.0075	-0.297*	0.523**	-0.0885	0.1223	0.294*
FLL	P					1	0.0998	0.290*	-0.2015	-0.1855	0.291*	-0.0250	-0.0557	0.2035
	G					1	0.1386	0.331*	0.358*	-0.280*	0.451**	0.0310	-0.1126	0.311*
FLW	P						1	0.1109	-0.0745	0.229	0.379**	0.1598	-0.0018	0.426**
	G						1	0.2250	-0.507**	0.650**	0.782**	0.2233	-0.1283	0.649**
S/P	P							1	-0.249*	0.324*	0.313*	0.1411	0.1317	0.346*
	G							1	-0.410**	0.602**	0.444**	0.339*	0.1614	0.459**
SF	P								1	0.3491*	-0.0726	-0.1851	0.258*	-0.2125
	G								1	-0.573**	-0.1442	-0.568**	0.345*	-0.424**
DM	P									1	0.1050	0.282*	-0.1330	0.305*
	G									1	0.2213	0.477**	-0.1486	0.412**
BY/H	P										1	-0.1532	0.0922	0.659**
	G										1	0.491**	0.1425	0.879**
HI	P											1	0.284*	0.638**
	G											1	0.526**	0.849**
TW	P												1	0.274*
	G												1	0.343*

DF50: Days to 50% Flowering, DM: Days to Maturity, PH: Plant Height, NT/H: Number of Tillers per Hill, NP/H: Number of Panicles per Hill, PL: Panicle Length, FLL: Flag Leaf Length, FLW: Flag leaf width, S/P: Spikelets per Panicle, DM: Days to Maturity, BY/H: Biological Yield per Hill, HI: Harvest index, TW: Test weight, P: Phenotypic, G: Genotypic

Table 6. Direct and indirect effects of yield component traits on grain yield in 25 Upland rice genotypes at both phenotypic and genotypic level

		DF50	PH	NT/H	NP/H	PL	FLL	FLW	S/P	SF	DM	BY/H	HI	TW	GY/H
DF50	P	0.0448	-0.0015	-0.0048	-0.0017	-0.0129	-0.0003	0.0087	0.0170	-0.0105	0.0227	0.0089	0.0073	-0.0010	0.311*
	G	0.0562	-0.0017	-0.0139	-0.0046	-0.0250	0.0029	0.0425	0.0507	-0.0365	0.0643	0.0099	0.0301	-0.0041	0.392*
PH	P	-0.0004	0.0119	0.0006	0.0021	0.0039	0.0051	0.0017	0.0022	-0.0025	0.0008	0.0035	0.0017	-0.0003	0.337*
	G	0.001	-0.0370	-0.0025	-0.0085	-0.0228	-0.0230	-0.0135	-0.0114	0.0199	-0.0065	-0.0268	-0.0026	0.0015	0.501*
NT/H	P	-0.0019	0.0008	0.0176	0.0149	0.0073	0.0017	0.0074	-0.0059	-0.0023	-0.0009	0.0034	0.0034	0.0009	0.312*
	G	0.0017	-0.0005	-0.0070	-0.0063	-0.0045	-0.0006	-0.0044	0.0025	0.0013	0.0013	-0.0020	-0.0017	-0.0003	0.333*
NP/H	P	-0.0003	0.0014	0.0068	0.0080	0.0036	0.0010	0.0043	-0.0021	-0.0010	-0.0001	0.0027	0.0021	0.0010	0.472*
	G	0.0056	-0.0157	-0.0616	-0.0682	-0.0515	-0.0096	-0.0550	0.0198	0.0170	0.0030	-0.0360	-0.0199	-0.0105	0.505*
PL	P	0.0021	-0.0024	-0.0030	-0.0032	-0.0072	-0.0013	-0.0022	0.0011	-0.0001	0.0013	-0.0015	0.000	-0.0006	0.1589
	G	-0.0771	0.1071	0.1122	0.1308	0.1732	0.0585	0.1026	-0.0201	0.0013	-0.0514	0.0906	-0.0153	0.0212	0.294*
FLL	P	0.0001	-0.0039	-0.0009	-0.0012	-0.0017	-0.0092	-0.0009	-0.0027	0.0018	0.0017	-0.0027	0.0002	0.0005	0.2035
	G	0.0037	0.0443	0.0061	0.0100	0.0241	0.0714	0.0099	0.0236	-0.0256	-0.0200	0.0322	0.0022	-0.0080	0.311*
FLW	P	0.0006	0.0004	0.0012	0.0016	0.0009	0.0003	0.0030	0.0003	-0.0002	0.0007	0.0011	0.0005	0.000	0.426**
	G	-0.0252	-0.0121	-0.0211	-0.0268	-0.0197	-0.0046	-0.0333	-0.0075	0.0169	-0.0216	-0.0260	-0.0074	0.0043	0.649**
S/P	P	-0.0017	-0.0008	0.0015	0.0012	0.0007	-0.0013	-0.0005	-0.0046	0.0011	-0.0015	-0.0014	-0.0006	-0.0006	0.346*
	G	-0.0621	-0.0212	0.0244	0.0200	0.0080	-0.0228	-0.0155	-0.0689	0.0282	-0.0415	-0.0306	-0.0233	-0.0111	0.459**
SF	P	0.0018	0.0016	0.0010	0.0010	-0.0001	0.0016	0.0006	0.0019	-0.0077	0.0027	0.0006	0.0014	-0.0020	-0.2125
	G	-0.0398	-0.0329	-0.0119	-0.0152	0.0005	-0.0219	-0.0311	-0.0251	0.0612	-0.0351	-0.0088	-0.0348	0.0211	-0.424**
DM	P	-0.0063	-0.0009	0.0007	0.0001	0.0022	0.0023	-0.0028	-0.0040	0.0044	-0.0125	-0.0013	-0.0035	0.0017	0.305*
	G	0.0814	0.0126	-0.0130	-0.0031	-0.0211	-0.0199	0.0462	0.0428	-0.0407	0.0711	0.0157	0.0339	-0.0106	0.412**
BY/H	P	0.1468	0.2208	0.1430	0.2522	0.1567	0.2156	0.2810	0.2319	-0.0539	0.0779	0.7419	-0.1137	0.0684	0.659**
	G	0.0824	0.3391	0.1361	0.2477	0.2450	0.2114	0.3664	0.2081	-0.0675	0.1037	0.4685	0.2299	0.0668	0.879**
HI	P	0.1190	0.1013	0.1418	0.1877	0.0022	-0.0182	0.1162	0.1029	-0.1350	0.2059	-0.1117	0.7292	0.2067	0.638**
	G	0.2917	0.0386	0.1357	0.1588	-0.0482	0.0169	0.1216	0.1846	-0.3094	0.2599	0.2673	0.5446	0.2865	0.849**
TW	P	0.0002	0.0002	-0.0004	-0.0009	-0.0006	0.0004	0.0000	-0.0010	-0.0019	-0.0010	-0.0007	-0.0021	-0.0074	0.274*
	G	0.0048	0.0027	-0.0032	-0.0103	-0.0082	0.0075	0.0086	-0.0108	-0.0231	0.0099	-0.0095	-0.0352	-0.0669	0.343*

DF50: Days to 50% Flowering, DM: Days to Maturity, PH: Plant Height, NT/H: Number of Tillers per Hill, NP/H: Number of Panicles per Hill, PL: Panicle Length, FLL: Flag Leaf Length, FLW: Flag leaf width, S/P: Spikelets per Panicle, DM: Days to Maturity, BY/H: Biological Yield per Hill, HI: Harvest Index, TW: Test weight, P: Phenotypic, G: Genotypic

4. CONCLUSION

From the present investigation, it is concluded that studied advanced breeding lines of upland rice showed significant variation to all the characters. On the basis of mean performance among 25 genotypes of upland rice, SHUATS UPR – 34 (4.52 tons/ha) was found to be superior in grain yield per hectare followed by SHUATS UPR – 33 (4.38 tons/ha) over the check variety SAHBHAGIDHAN (4.37 tons/ha). Estimation of genetic variability parameters, correlation analysis and path coefficient analysis revealed that, number of tillers per hill, number of panicles per hill, days to 50% flowering, plant height, biological yield per hill were most reliable traits for yield improvement. So, the utmost importance should be given to these characters during the selection for yield improvement in upland rice.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Biswas CD, Biswas BK, Sadat Md, Saha PS, Tareq Md. Morphological comparison among upland and lowland rice varieties for development of drought resistant variety. *Int. J. Bus. Soc. Sci. Res.* 2018; 6(4):96-102.
2. Barker R, Dawe D, Tuong T, Bhuiyan S, Guerra L. The outlook for water resources in the year 2020: challenges for research on water management in rice production. *Southeast Asia.* 1999;1:1-5.
3. Tuong TP, Bouman BAM. Rice production in water-scarce environments, CABI Publishing, Wallingford. 2003;53-67.
4. Bouman BAM, Lampayan RM, Tuong TP. Water management in irrigated rice: coping with water scarcity. International Rice Research Institute, Philippines. 2007;54.
5. Romana GS. Direct seeded rice versus normal transplanted rice: An economic comparison. *Indian J. Econ. Dev.* 2014; 10(2):117.
6. Ebadi A, Mommadi SA, Sadeghzadeh B, Kamrani M. Study of heritability and genetic advance of agronomic traits in Barley (*Hordeum vulgare* L.) and graphic analysis of trait relations by Biplot. *Jordan J. Biol. Sci.* 2016;12(1):299-310.
7. Ayer DK, Sharma A, Ojha BR, Paudel A, Dhakal K. Correlation and path coefficient analysis in advanced wheat genotypes. *SAARC J. Agri.* 2017;15(1):1-12.
8. Wamanrao AN, Kumar V, Meshram D. Correlation and path coefficient analysis of grain yield and its growth components in soybean (*Glycine max.* L.). *Int. J. Curr. Microbiol. App. Sci.* 2020;9(3):2445-2451.
9. Swapnil, Prasad K, Chakraborty M, Singh DN, Ekka JP, Kumari P. Genetic variability, correlation and path coefficient studies in F3 population of rice (*Oryza sativa* L.). *Int. J. Curr. Microbiol. App. Sci.* 2020;9(9): 1304-1312.
10. Fisher RA. XV. The correlation between relatives on the supposition of Mendelian inheritance. *Trans R Soc Edinb.* 1919; 52(2):399-433.
11. Burton GW. Quantitative inheritance in grasses. *Proceedings of the 6th intern grasslands congress.* 1952;1:277-83.
12. Burton GW, Devane EH. Estimating heritability in tall fescue (*Festuca Arundinaceae*) from replicated clonal material. *Agron J.* 1953;51:515-8.
13. Johnson HW, Robinson HF, Comstock RE. Estimation of genetic and environmental variability in Soyabean. *Agron J.* 1955; 47(7):314-8.
14. Al-Jibouri HA, Miller PA, Robinson HF. Genotypic and environmental variation and correlation in upland cotton cross of interspecies origin. *Agron J.* 1958; 50(10):633-6.
15. Fisher RA, Yates F. Statistical tables for biological, agricultural and medical research. 6th Ed. Oliver and Boyd, Edinburgh and London. 1963;146.
16. Girma BD, Kitil MA, Banje DG, Biru HM, Serbessa TB. Genetic variability study of yield and yield related traits in rice (*Oryza sativa* L.) genotypes. *Adv. Crop. Sci. Tech.* 2018;6(4):2329-8863.
17. Magar BT, Acharya S, Gyawali B, Timilsena K, Upadhayaya J, Shrestha J. Genetic variability and trait association in maize (*Zea mays* L.) varieties for growth and yield traits. *Heliyon.* 2021;e07939.
18. Kumar S, Chauhan MP, Tomar A, Kasana RK. Coefficient of variation (GCV&PCV), heritability and genetic advance analysis for yield contributing characters in rice (*Oryza sativa* L.). *Int. J. Pharmacognosy. Phytochem. Res.* 2018;7(3):2161-2164.
19. Khatun MST, Hanfi MM, Yusop MR, Wong MY, Salleh FM, Ferdous J. Genetic variation, heritability, and diversity analysis of upland rice (*Oryza sativa* L.) genotypes

- based on quantitative traits. Biomed Res. Int. 2015;290861.
20. Chandra B, Reddy T, Sudheer S. Variability parameters for yield, its components and quality traits in rice (*Oryza sativa* L.). Crop Res. 2009;38(1,2&3):146-146.
 21. Anjaneyulu M, Reddy DR, Reddy KHP. Genetic variability, heritability and genetic advance in rice (*Oryza sativa* L.). Res. Crops. 2010;11(2):415-416.
 22. Lush JL. Inter-size correlation regression of offspring on dairy as a method of estimating heritability of characters. Proceedings of the American Society of Animal Production. 1940;33:293-301.
 23. Tiwari GC. Variability, heritability and genetic advance analysis for grain yield in rice. Int. J. Eng. Res. Appl. 2015;(5):46-49.
 24. Pratap A, Bisen P, Loitongbam B, Singh PK. Assessment of genetic variability for yield and yield components in rice (*Oryza sativa* L.) germplasm. Int. J. Bio-Resour. Stress manag. 2018;9(1):87-92.
 25. Longjam S, Singh NB. Assessment of heritability and genetic advance for yield contributing characters in hill rice (*Oryza sativa* L.) genotypes of Manipur. J. Pharma. Innov. 2019;8(4): 07-11.
 26. Singh VK, Wahi N, Mishra SK, Singh BD, Singh NK. Studies on genetic variability, correlation analysis, character association and path analysis of phenotypic characteristics of twelve mega varieties of rice and its near-isogenic lines carrying high grain number per panicle QTL qGN4. Curr. Trends. Biotechnol. Pharm. 2022; 16(1):35-45.
 27. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in soybeans. Agron. J. 1955; 47:314-318.
 28. Rashmi D, Saha S, Loitongbam B, Singh S, Singh PK. Genetic variability study for yield and yield components in rice (*Oryza sativa* L.). Int. J. Agric. Environ. Biotechnol. 2017;10(2):171.
 29. Singh R, Yadav V, Mishra DN, Yadav A. Correlation and path analysis studies in rice (*Oryza sativa* L.). Int. J. Pharmacognosy. Phytochem. Res. 2018; 2084-2090.
 30. Srijan A, Kumar SS, Raju CH, Jagadeeshwar R. Character association and path coefficient analysis for grain yield of parents and hybrids in rice (*Oryza sativa* L.). J. Appl. Nat. Sci. 2016;8(1):167-172.
 31. Wright S. Correlation and causation. J. Agric. Res. 1921;20:557-85.
 32. Sharma MK, Sharma AK. Character association and path analysis of yield and its components in direct seeded upland rice (*Oryza sativa* L.). Progress. Agric, 2009;9(1):117-120.
 33. Mishu FM, Rahman W, Azad MAK, Biswas BK, Talukder AI, Kayers O, Islam R, Alam R. Study on genetic variability and character association of aromatic rice (*Oryza sativa* L.) cultivar. Int. J. Plant Soil Sci. 2016;9(1):1-8.
 34. Belete DA, Tewachew A, Mulugeta B, Mulualem T. Correlation and path coefficient studies for yield and its components of upland rice (*Oryza sativa* L.) in north western Ethiopia. J. Sci. Agric. 2022;6:14-19.

© 2022 Rajasekhar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92570>