



Damage Control Resuscitation: Evidence-based Systematic Review for Treating Traumatized Patients

Lucas Pessoa Souza ^{a++},
Pedro Figueiredo Martins Ribeir ^{a++},
Roberto Cesar Lima Santos ^{a++},
Rodolfo Feitosa de Figueiredo Moreira ^{a++},
Petrus Moura Andrade Lima ^{b#}
and Leslie Clifford Noronha Araujo ^{at*}

^a UNICAP (Catholic University of Pernambuco), Brazil.

^b SES (State Health Secretary of Pernambuco), Brazil.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jammr/2024/v36i65477>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117759>

Original Research Article

Received: 06/05/2024

Accepted: 07/06/2024

Published: 10/06/2024

ABSTRACT

Aim: The aim of this study is to analyze concepts and updates on Damage Control Resuscitation (DCR), focusing on critical patient approaches, therapeutic failures identification, and the relevance of damage control in trauma.

⁺⁺ Medical Student;

[#] Physician, General Surgeon;

[†] Adjunct Professor;

^{*}Corresponding author: E-mail: leslie.clifford@gmail.com;

Cite as: Souza , Lucas Pessoa, Pedro Figueiredo Martins Ribeir, Roberto Cesar Lima Santos, Rodolfo Feitosa de Figueiredo Moreira, Petrus Moura Andrade Lima, and Leslie Clifford Noronha Araujo. 2024. "Damage Control Resuscitation: Evidence-Based Systematic Review for Treating Traumatized Patients". *Journal of Advances in Medicine and Medical Research* 36 (6):343-49. <https://doi.org/10.9734/jammr/2024/v36i65477>.

Methods: This article presents a systematic review of damage control resuscitation in trauma, encompassing both consolidated and emerging literature, using informative tables in accordance with PRISMA 2020 guidelines. Conducted between March 2023 and March 2024, using the research descriptor "Resuscitation for damage control" in PubMed. Studies published between 2016 and 2023, freely accessible in English and Portuguese, were included, while those not aligning with the objective were excluded. Out of 979 articles found, 10 were selected after applying inclusion and exclusion criteria.

Results: Damage Control Resuscitation (DCR) is based on three pillars: permissive hypotension, minimization of crystalloid use, and balanced resuscitation. This approach aims to maintain adequate blood pressure, limit crystalloid use to avoid complications, and balance blood product transfusion. Hypocalcemia has emerged as a new challenge in the lethal triad of trauma, impacting coagulation and mortality. New techniques, such as REBOA and viscoelastic tests (TEG and ROTEM), are being explored to improve resuscitation in severe trauma cases. Tranexamic acid (TXA) also stands out as an effective treatment for excessive bleeding in trauma situations. The importance of pre-hospital care and early interventions is emphasized to optimize bleeding control and resuscitation in traumatized patients.

Conclusion: Damage Control Resuscitation (DCR) is an evidence-based strategy for treating traumatized patients, focusing on permissive hypotension, reduced crystalloid use, and balanced transfusion. In resource-limited settings, alternatives such as whole blood use can be adopted. Pre-hospital care and early interventions are crucial for better outcomes and survival.

Keywords: Damage control; hypovolemic shock; traumatized patients; resuscitation.

1. INTRODUCTION

The Resurrection Damage Control (RCD) technique originated in the military context, during the Vietnam War (1955-1975), when it was instructed that seriously injured soldiers often died during transport to the hospital due to loss of limb. Weight, blood flow and impairment of cardiovascular function. Faced with this situation, military doctors developed a temporary surgical technique that consists of carrying out limited surgical exploration, temporarily controlling the most critical injuries and stabilizing the patient before starting definitive treatment. This technique is called "damage control resuscitation" [1].

The DCR is an eligible protocol in many scenarios, whether emergency room, intensive care unit, operating room and/or interventional radiology. Patients with severe shock associated with imminent metabolic repercussions and multiple organ injuries, which would require a long time for definitive correction. The literature taught that searching for a definitive approach in critically unstable patients often leads to death during initial care or a few hours later [2-4].

The demand to cover these patients with a severe profile, the need in favor of the rapid control of hemodynamic instability caused by trauma emerged from the DCR protocol, being

used around the world. The frequent updating of the protocol and the controversial pillars of the DCR, became mandatory to the healthcare team periodically review all protocols, especially those providing first care, in order to avoid avoidable fatal outcomes.

The aim of the study is to analyze concepts and updates on DCR, focusing on critical patient approaches, therapeutic failures identification, and the relevance of damage control in trauma [4,5].

2. METHODOLOGY

Searched between March 2023 and April 2024 from the PubMed, Scielo and Lilacs database, using the descriptor in Health Sciences (Decs): "Resuscitation for damage control" selecting full access Papers provided for research by the Catholic University of Pernambuco (UNICAP) which fulfill the inclusion criteria.

The inclusion criteria were papers published between 2016 and 2023 in English or Portuguese presented a consistent with the objective of the research, and using the descriptor above. After applying the criteria, from 979 papers in primary search, 10 were selected, read and summarized. The selection of articles in this particular period aimed to follow the most recent evidence of scientific production on the topic. As an exclusion criterion, articles that did

not respond to the research object or having less based evidence, such as case reports.

3. RESULTS

According to the information obtained in the reviewed studies, the management of DCR is based on three important pillars, permissive hypotension, minimization of the use of crystalloids and balanced resuscitation. The first principle brings an approach to blood pressure management aimed at ensuring adequate perfusion of organs, avoiding exacerbation of bleeding. The strategy involves maintaining a "hypotensive" state with a target systolic blood pressure of 80 mmHg (or a mean arterial pressure of 80 mmHg in cases of neurotrauma)

to minimize future blood loss and prevent displacement of blood clots [6].

The reduction of using crystalloids, emerged as one of the three principles of RCD. Previously, liberal use of these fluids was associated with serious complications, including coagulopathy, hypothermia, and acidosis [7].

The third pillar of DCR, we can emphasize the use of balanced proportions of transfusion of blood products, particularly plasma, platelets and red blood cells (RBCs), in a ratio of 1:1:1, approaching whole blood [7-9].

The Table 1 summarizes the following variables are shown: Title of the article, Authors/year and Proposed treatment.

Table 1. Papers synthesis

Title	Author, year	Porposal
Resuscitation for Hypovolemic Shock	Kyle J Kalkwarf, 2017.	Limit the use of large crystalloid transfusions. Ratio in resuscitation 1:1:1. Whole blood transfusion as superior method in resuscitation. Avoid hypothermia.
Reboa as a new damage control component in hemodynamically unstable noncompressible torso hemorrhage patients	Ordoñez CA, 2020	Propose two new deployment algorithms for hemodynamically unstable noncompressible torso hemorrhage patients: Blunt and REBOA
Ressuscitação para controle de danos	Jason M Samuels, 2017.	Use of pharmacological therapies designed to inhibit clot degradation in seriously injured patients, using tranexamic acid.
Damage control: Concept and implementation	Malgras, 2017	Permissive Hypotension, maintaining a systolic blood pressure around 80 mmHg.
Prehospital Damage Control: The Management of Volume, Temperature... and Bleeding!	Lugo, 2020	Consensus and summarizes the experience of the Trauma and Emergency Surgery Group in managing trauma intensive care over the past 30 years.
The Role of TEG and ROTEM in Damage Control Resuscitation	Jason B. Brill, 2021.	TEG and ROTEM can predict massive transfusion and development of coagulopathy
Whole blood for blood loss: hemostatic resuscitation in damage control	Juan Carlos Salamea-Molina, 2020.	Use of whole blood in resuscitation providing a practical alternative in resource-limited settings.
Optimal Fluid Therapy for Traumatic Hemorrhagic Shock	Chang, 2017.	Discusses the paradigm shift in traumatic hemorrhagic shock resuscitation with the advent of DCR
Traumatic hemorrhage and chain of survival	Latif RK, 2023.	The prehospital and hospital chain of survivorship care

4. DISCUSSION

The Letahl triad of trauma conditions may result in respiratory failure, cardiac arrhythmias, proteolysis and other circulatory dysfunctions with high lethal potential. Surgical patients who develop metabolic acidosis, especially hyperchloremic acidosis, have worse prognoses, both immediate and late. Also linked to acidosis, coagulopathy will further increase the patient's chances of mortality, given the reduced activity of coagulation factors and their complexes. Finally, hypothermia alters coagulability by reducing platelet enzymatic activity and the fibrinolytic system, thus completing the fatal cycle [10,11].

The literature shows the three pillars of DCR are a well-established strategy, the first principle brings an approach to blood pressure management aimed at ensuring adequate perfusion of organs, avoiding exacerbation of bleeding. The strategy involves maintaining a "hypotensive" state with a target systolic blood pressure of 80 mmHg (or a mean arterial pressure of 80 mmHg in cases of neurotrauma) to minimize future blood loss and prevent displacement of blood clots[3–7,9,12]. Placing in a unanimous position among the reviewed authors and studies.

Improved survival in trauma patients was identified with the reduction in crystalloid use, establishing this practice as the second pillar of DCR [7].

Previously, liberal use of these fluids was associated with serious complications, including coagulopathy, hypothermia, and acidosis. Crystalloids dilute clotting factors, cool the patient, cause acidosis, and can lead to edema and organ dysfunction. Furthermore, they are associated with a series of complications, such as heart, respiratory, gastrointestinal and immunological problems, increasing mortality. Limiting the use of crystalloids also reduces the need for transfusions in patients with severe injuries. [2,7,8,11].

Several authors have established as the last pillar the use of balanced proportions of transfusion of blood products, particularly plasma, platelets and red blood cells (RBCs), in the ratio of 1:1:1, since this proportion is close to whole blood.

Plasma transfusions are beneficial in reducing inflammation, edema and vascular permeability,

improving platelet function and alleviating blood hypercoagulability[8]. To speed up plasma transfusion, pre-thawed or liquid plasma is increasingly used, facilitating earlier transfusions and better results. Its implementation in the RCD led to a reduction in the use of crystalloids, morbidity from abdominal compartment syndrome, infection, organ failure and mortality from hemorrhage[7,8,13].

In environments with limited resources, where this practice may not be available, the use of whole blood appears as an alternative, having advantages of a physiological proportion of components, including red blood cells, plasma, platelets and clotting factors, which can help to restore the balance of these components in the patient's body. It may also be more convenient, as component therapy eliminates the need for separate storage and administration of different parts of the blood [14].

Ordoñez et al. in 2020, they noticed the presence of hypokalemia in 50% of patients before the start of blood transfusion, and suggested the inclusion of this condition as a new pillar to be corrected in polytraumatized patients. But there is still no consensus in the literature[15].

This state of pre-existing trauma is potentially worsened by initial treatment using DCR in which blood and blood products are transfused. Early management of hypocalcemia in trauma patients may be more significant than previously believed because as serum calcium levels fall, a state of coagulopathy occurs, which can lead to continued hemorrhage and possible death. However, there is still no well-established pathophysiology and management, further studies are still required to establish a better targeted approach[15,16].

Resuscitation efforts in patients requiring massive transfusion in trauma have been discussed, including viscoelastic tests. The importance of recognizing hypocoagulable tracings and promptly addressing modifiable factors that contribute to trauma-induced coagulopathy (TIC). Values obtained from viscoelastic testing help determine appropriate interventions such as plasma transfusion, anticoagulant reversal, and targeted treatments for anticoagulant-associated hemorrhage. Furthermore, the role of assessing fibrinolysis in guiding antifibrinolytic therapy, in parallel to the usually already established methods[11,13].

In therapy guided by viscoelasticity, it is possible to analyze the slow kinetics of the clot, such as its strength, thus identifying the moment or need to perform a fibrinogen or platelet transfusion. Although some studies suggest potential benefits of viscoelastic-guided resuscitation, more research is needed to establish its superiority over conventional approaches [13].

The tranexamic acid (TXA), which is a medicine used to prevent and treat excessive bleeding, acts to delay and decompose blood clots, thus reducing bleeding. TXA is particularly useful in situations where there is uncontrolled bleeding, such as trauma, surgery, heavy menstrual bleeding, or certain medical conditions such as hemophilia. One of the significant advances in the use of TXA came from the CRASH-2 trial, a large-scale study that demonstrated its effectiveness in reducing mortality in trauma patients when administered shortly after injury within 3 hours [17]. Since then, TXA has become a crucial component of trauma care protocols around the world. It is important to note that although TXA can be highly beneficial in certain situations, it is not suitable for everyone, and medical professionals must carefully assess each patient's condition before administering it [3].

Ordoñez and Cols, 2020 began to advocate in favor of Resuscitative Endovascular Balloon Occlusion of the Aorta (REBOA), as a new pillar in the control of hypovolemia [14,15]. In addition to the fact this treatment model needs a complex treatment center, a high qualified staff must be present, difficulting the implementation in many centers. Maybe for those reasons, the REBOA has not yet been established by other authors.

Despite the conduct already established in relation to DCR, the importance of pre-hospital care as the first step to guarantee bleeding control and resuscitation in traumatized patients must be reinforced. Furthermore, early interventions without increasing hospital transfer time are essential to optimize the success rate [18-20].

5. CONCLUSION

The multiple trauma patient, whether in the hospital or pre-hospital area, continues to be a challenge for the healthcare team, where early treatment prevents fatal outcomes, mainly

related to the lethal triad. Systematic action, especially in the early stages, improves results.

The Blood Component Resuscitation approach emerges as a multifaceted, evidence-based strategy for the effective management of trauma patients during the DCR. Through its three pillars - permissive hypotension, minimization of crystalloid use and balanced resuscitation.

The context of limited resources in some healthcare settings may require viable alternatives, such as the use of whole blood. Furthermore, the continued evolution of therapy, including use of viscoelastic testing, highlights the importance of adapting to each patient's individual needs. Tranexamic acid also stands out as a crucial tool in reducing excessive bleeding and the importance of a pre-hospital care early management.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ball C. Damage control resuscitation: history, theory and technique. *Canadian Journal of Surgery* 2014;57:55–60. Available: <https://doi.org/10.1503/cjs.020312>.
2. Stone HH, Strom PR, Mullins RJ. Management of the Major Coagulopathy with Onset during Laparotomy. *Ann Surg* 1983;197:532–5. Available: <https://doi.org/10.1097/0000658-198305000-00005>.

3. Samuels JM, Moore HB, Moore EE. Damage Control Resuscitation. *Chirurgia (Bucur)* 2017;112:514.
Available:<https://doi.org/10.21614/chirurgia.112.5.514>.
4. González Posada MA, Biarnés Suñe A, Naya Sieiro JM, Salvadores de Arzuaga CI, Colomina Soler MJ. Damage control resuscitation in polytrauma patient. *Rev Esp Anestesiología y Reanimación*. 2019;66.
Available:<https://doi.org/10.1016/j.redar.2019.03.009>.
5. Giannoudi M, Harwood P. Damage control resuscitation: lessons learned. *European Journal of Trauma and Emergency Surgery*. 2016;42.
Available:<https://doi.org/10.1007/s00068-015-0628-3>
6. Malgras B, Prunet B, Lesaffre X, Boddart G, Travers S, Cungi PJ, et al. Damage control: Concept and implementation. *J Visc Surg*. 2017;154.
Available:<https://doi.org/10.1016/j.jviscsurg.2017.08.012>.
7. Kalkwarf KJ, Cotton BA. Resuscitation for hypovolemic shock. *Surgical Clinics of North America*. 2017;97:1307–21.
Available:<https://doi.org/10.1016/j.suc.2017.07.011>.
8. Chang R, Holcomb JB. Optimal Fluid Therapy for Traumatic Hemorrhagic Shock. *Crit Care Clin*. 2017;33.
Available:<https://doi.org/10.1016/j.ccc.2016.08.007>.
9. Chang R, Eastridge BJ, Holcomb JB. Remote damage control resuscitation in austere environments. *Wilderness Environ Med*. 2017;28.
Available:<https://doi.org/10.1016/j.wem.2017.02.002>.
10. Silva Junior JM da. Acidose metabólica em pacientes cirúrgicos de alto risco: Importância prognóstica. Universidade de São Paulo; 2016.
Available:<https://doi.org/10.11606/T.5.2016.tde-03022016-154257>.
11. Nascimento Júnior B, Scarpelini S, Rizoli S. Coagulopatia no trauma. *Medicina (Ribeirão Preto)*. 2007;40:509–17.
Available:<https://doi.org/10.11606/issn.2176-7262.v40i4p509-517>.
12. Lagrone LN, Stein D, Cribari C, Kaups K, Harris C, Miller AN, et al. American association for the surgery of Trauma/American college of surgeons committee on trauma: Clinical protocol for damage-control resuscitation for the adult trauma patient. *Journal of Trauma and Acute Care Surgery*. 2024;96.
Available:<https://doi.org/10.1097/TA.0000000000004088>.
13. Brill JB, Brenner M, Duchesne J, Roberts D, Ferrada P, Horer T, et al. The Role of TEG and ROTEM in Damage Control Resuscitation. *Shock*. 2021;56.
Available:<https://doi.org/10.1097/SHK.0000000000001686>.
14. Ordoñez CA, Parra MW, Caicedo Y, Padilla N, Rodríguez-Holguín F, Serna JJ, et al. Reboa as a new damage control component in hemodynamically unstable noncompressible torso hemorrhage patients. *Colomb Med*. 2020;51.
Available:<https://doi.org/10.25100/cm.v51i4.4506>.
15. Ordoñez CA, Parra MW, Serna JJ, Rodríguez-Holguín F, García A, Salcedo A, Damage control resuscitation: Reboa as the new fourth pillar. *Colomb Med*. 2020;51.
Available:<https://doi.org/10.25100/cm.v51i4.4353>.
16. Salamea-Molina JC, Himmler AN, Angel LIV, Ordoñez CA, Parra MW, Caicedo Y, et al. Whole blood for blood loss: hemostatic resuscitation in damage control. *Colomb Med*. 2020;51.
Available:<https://doi.org/10.25100/cm.v51i4.4511>.
17. Latif RK, Clifford SP, Baker JA, Lenhardt R, Haq MZ, Huang J, et al. Traumatic hemorrhage and chain of survival. *Scand J Trauma Resusc Emerg Med* 2023;31.
Available:<https://doi.org/10.1186/s13049-023-01088-8>.
18. Meléndez-Lugo JJ, Caicedo Y, Guzmán-Rodríguez M, Serna JJ, Ordoñez J, Angamarca E, et al. Prehospital Damage Control: The Management of Volume, Temperature... and Bleeding! *Colomb Med*. 2020;51.
Available:<https://doi.org/10.25100/cm.v51i4.4486>.
19. Innab A, Bahari G. Post-Traumatic Stress Disorder and Associated Factors among Caregivers of Patients with COVID-19 in Saudi Arabia: A Cross-sectional

Study, Journal of Pharmaceutical Research International. 2022;34(13A): 33–40.
DOI: 10.9734/jpri/2022/v34i13A35572

20. Giannoudi M, Harwood P. Damage control resuscitation: lessons learned. European Journal of Trauma and Emergency Surgery. 2016;42:273-82.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/117759>