



Review on Pharmacological Profile of Medicinal Vine: *Tinospora cordifolia*

Kumar Antul^{1*}, Paul Amandeep¹, Singh Gurwinder¹ and Choudhary Anuj¹

¹Department of Botany, College of Basic Sciences and Humanities, Punjab Agricultural University,
Ludhiana-141004, Punjab, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2019/v35i530196

Editor(s):

(1) Dr. Yahya Elshimali, Professor, Department of Internal Medicine, Charles Drew University of Medicine and Science, USA.

Reviewers:

(1) Mustafa Sevindik, Akdeniz University, Turkey.

(2) Senthil Kumar Raju, Swamy Vivekanandha College of Pharmacy, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/49114>

Review Article

Received 03 March 2019

Accepted 19 May 2019

Published 03 June 2019

ABSTRACT

Tinospora is highly distributed in the tropical and subtropical region of India. This climbing deciduous shrub widely reported in China, Bangladesh and Srilanka. The plant is rich in many phytoconstituents that are useful in drug designing. It is highly used against cancer, tumour suppression, and act as an anti-allergic compound. It is commonly known as gudhuchi, belongs to the family Menispermaceae. *Tinospora* is most valuable herb known for its medicinal properties from Vedic periods and cures various diseases such as malaria, asthma and urinary disorders. The genus *Tinospora* consists of many classes of chemicals such as alkaloids, diterpenoids lactones, steroids, aliphatic compounds and polysaccharides. It is the best remedy for both children as well as adults against respiratory tract diseases. The plant shows various antioxidant, anti-hyperglycemic, anti-neoplastic and hepatoprotective properties. In this review article medicinal property, chemical constituents and full description have been explored.

Keywords: *Tinospora*; gudhuchi; diterpenoids; antioxidant; anti-neoplastic; steroids.

*Corresponding author: E-mail: antulkumar007@gmail.com;

1. INTRODUCTION

Guduchi or *Giloya* is the most commonly used plant which contains a large number of valued products. It has a wide history in the Indian medicinal system and considered one of the best *Rasayana* and is unusual in its potent versatility. In recent years, significant progress has been attained for its biological activity and medicinal applications. It is a semi-woody climbing shrub that is deciduous and perennial. This herbaceous vine grows on hedges and trees described as “one which protects the body”. It is often seen growing up Mango or Neem trees. Herbalist Sebastian Pole writes that “those growing up neem trees are said to be the best as the synergy between these two bitter plants enhances *guduchi*'s efficacy.” It is indigenous to areas of India, Myanmar, and Sri Lanka [1, 36]. *Guduchi* typically grows in deciduous and dry forests at elevations up to 1000 ft. The leaves are heart-shaped (*cordifolia*) and mucilaginous. Its stems, when fresh, have a green succulent bark covered by a thin brown bark and are studded with warty lenticels. When dry, the stem shrinks and the bark separate from the wood. The roots are long narrow aerial roots that arise from the branches [36]. The stems, leaves, and roots are used in medicine. All three parts should be collected in the summer when the bitter qualities are most abundant and, if not used fresh, dried in the shade. *Guduchi* grows well without fertilizer or pesticide making it simple to grow. It is easy to recognize and can be propagated by cuttings. *Guduchi* is a large glabrous deciduous climbing shrub belonging to the family Menispermaceae [37]. It is distributed throughout tropical Indian subcontinent and China, ascending to an altitude of 300 m. In Hindi, the plant is commonly known as *Giloya* or *Amrita* which is a Hindu mythological term that refers to the heavenly elixir that has saved celestial beings from old age and kept them externally young [2]. The stem of *T. cordifolia* is rather succulent with long filiform fleshy aerial roots from the branches. The bark is creamy white to grey, deeply left spirally, the space in between being spotted with large rosette-like lenticel. The leaves are membranous and cordate. The flowers are small and yellow or greenish yellow [3]. In auxiliary and terminal racemes or racemose panicles, the male flowers are clustered and female is usually solitary. The drupes are ovoid, glossy, succulent, red and pea-sized. The seeds are curved and pea-sized [4]. Fruits are pear-shaped, fleshy, shiny turn red when boiled. *Guduchi* is used as a *Rasayana*

due to its potency of enhancing longevity and vitality. It is widely used in ayurvedic for a variety of purposes associated with inflammation allergies, neurology and glucose metabolism, general debility, fever, diabetes, dyspepsia, urinary infection, jaundice and skin diseases. In today's world of modern medicine, it is also called a magical herb due to its property to treat a lot of diseases.

2. CLASSIFICATION

Kingdom : Plantae
 Division : Magnoliophyta
 Class : Magnoliopsida
 Order : Ranunculales
 Family : Menispermaceae
 Genus : *Tinospora*
 Species : *cordifolia*

2.1 Total Species

Some observers found that there are total 15 species and out of these some of the medicinally important species are *T. cordifolia*, *T. crispa*, *T. cordifoli*, *T. malabarica*, *T. tomentosa*, *T. uliginosa* etc [1].

2.2 Vernacular Names

Assamese : *Siddhilata, Amaralata*
 Bengali : *Gulanchara*
 English : *Heartleaf moonseed*
 Gujarati : *Galac, Garo*
 Hindi : *Giloe, Gurchara*
 Kannada : *Amrutaballi*
 Kashmiri : *Amrita, Gilo*
 Malayalam : *Chittamrutu*
 Marathi : *Gulvel*
 Oriya : *Guluchi*
 Punjabi : *Gilo*
 Sanskrit : *Amrit*
 Tamil : *Seendal, Seendi Kodi*
 Telugu : *Thippateega*
 Urdu : *Abb-e-Hyat*

3. BOTANICAL DESCRIPTION

T. cordifolia is a large, perennial, deciduous, climbing shrub with the succulent stem. The stem is fibrous and having wedge-shaped wood bundles with large vessels. The bark is papery, creamish white in colour, left spirally and stem containing rosette-like lenticles. The leaves are simple, alternate and cordate in shape also consist of 7-9 nerves on the entire leaf [5]. Flowers are axillary, small, cymose, yellow-greenish in colour. Male and female flowers are

always originated on separate branches. Male flowers are present in cluster form while female flowers are in the solitary form [6]. The best time for the growth of flower is during summer [7]. Sometimes small yellowish flowers are also present on long spikes. Fruits of *Tinospora* are pea-shaped shiny, draping and become red when fully grown. Fruits are generally single-seeded and fleshy. The fruits get maturity in the winter season. Seeds are hooked or curved in shape. The root portion is aerial, thread-like, long, fleshy and is in branching form.

4. HABITAT AND DISTRIBUTION

T. cordifolia prefers subtropical and tropical for growth. For better cultivation, light-medium sandy loam soil rich in organic matter and with adequate drainage is suitable. This plant is highly grown tropical India, South Asia, Indonesia, Philippines, Thailand and China. The plant is also observed from the South East Asian continent such as Malaysia, Indonesia and Tamilnadu.

5. CLIMATE AND SOIL

The plants preferred subtropical and tropical conditions for proper growth. For better cultivation, light-medium sandy loam soil rich in organic matter and with adequate drainage is suitable. It shows low resistance towards high rainfall or waterlogged conditions. Stem cutting is the best method to enhance commercial use.

6. FLORAL AND FRUIT STUDY

Inflorescence starts in the summer season. The male flowers are small in size, yellow or green in colour, and occur in groups. While female flowers are usually green and solitary in nature. The fruit size and shape is like a pea pod and turns green to red when ripe in winter.

7. CHEMICAL CONSTITUENTS

A number of chemical constituents have been extracted from the different parts of *T. cordifolia*. These chemical constituents belong to different classes viz; alkaloids, diterpenoid lactones, steroids, glycosides aliphatic compounds, polysaccharides. The main constituents of this plant are tinosporone, tinosporic acid, cordifolisides A to E, syringe, berberine, gilonin [8]. The chemical constituents in different parts and its uses are given in the Table 2.

It also contains various other chemicals like flavonoids, glycosides, saponins and little amount of phytosterols. These chemicals show antioxidant activity when these are combined with other drugs. These main constituents which are present in a very high amount are alkaloids and terpenes. The leaves are the rich source of protein, calcium as well as phosphorus.

8. TOXICOLOGY

In human beings, toxic effects of *Tinospora* is very less known. But sometimes its high dose causes some harmful effects on the body. It might lower blood sugar level, use it carefully if anyone has diabetes, it also increases the symptoms of autoimmune symptoms. It is also advised to avoid the intake of *Tinospora* during pregnancy and breastfeeding time [16].

9. AYURVEDIC PROPERTIES

Ayurveda is one of the most ancient medical sciences of the world. Rasayana is one of the eight clinical specialities of classical Ayurveda [37]. The concept of Rasayana therapy is not a single drug. Treatment but it is a comprehensive and specialized regimen capable of producing healthful longevity and improved mental facilities. Several medicinal plants have been described as Rasayana in Ayurveda [17].

Table 1. Different part used and its benefits

Part	Part used	Benefits
1.	Leaves	Juice or decoction of leaves is taken orally with honey in case of fever.
2.	Whole plant	Anti-pyretic
3.	Roots	The roots are used as antidote to snake bite and scorpion sting after combining with other drugs.
4.	Stem	It is bitter in taste, stimulates bile secretion, stomachic, diuretic, removing burning sensations, vomiting and also cure jaundice.




Fig. 1. Schematic representation of different parts of *Tinospora* plant and their folk uses




Fig. 2. Pharmacological property of *Tinospora cordifolia*

Table 2. Plant part used, chemical constituents and effect on humans

Sr. No.	Part	Chemical constituents	Uses	References
1.	Used Whole Plant	β -sitosterol, δ -sitosterol, 20- β -Hydroxy ecdysone, Furanolactone, Clerodane derivatives and [(5R,10R)-4R-8Rdihydroxy- 2S-3R:15,16-diepoxycleroda- 13 (16), 14-dieno-17,12S:18,1S-dilactone] and Tinosporon, Tinosporides, and Jateorine, Columbin, Octacosanol, Heptacosanol, Miscellaneous- Nonacosan-15-one3, (α ,4-di hydroxy- 3-methoxy-benzyl)-4-(4- hydroxy-3-methoxy-benzyl)-tetrahydrofuran, Tinosporidine, Cordifol, Cordifellone, N-transferuloyl tyramine as diacetate, Giloin, Giloinin, Tinosporic acid.	<ul style="list-style-type: none"> ➤ Anti-stress activity ➤ Antidote to snake bite and scorpion sting, ➤ Analgesic and neuro-pharmacological activities, Diabetes, Rheumatoid arthritis, Gout, cancer, high cholesterol content Anti-asthmatic and chronic cough treatment, ➤ Antipyretic and anti-inflammatory activity, Anaemia, jaundice, normalization of altered liver function, ➤ Cardiac disorder, ➤ Anti-leprotic, ➤ Gastrointestinal and anti-ulcer activity ➤ Anti-fertility activity, Hepatoprotective activity 	[9] [10] [11]
2.	Root	3, (a,4-di hydroxy-3-methoxy-benzyl)-4-(4- compounds hydroxy-3-methoxy-benzyl)- tetrahydrofuran, Jatrorrhizine, Tinosporidine, Cordifol, Cordifellone, Giloinin, Giloin, N-transferuloyl tyramine as diacetate, Tinosporic acid. Berberine, Choline, Tembetarine, Magnoflorine, Tinosporin, Palmetine, Isocolumbin, Aporphine alkaloids, Jatrorrhizine, Tetrahydropalmatine, sitosterol	<ul style="list-style-type: none"> ➤ Anti-neoplastic ➤ Anti-oxidant ➤ Anti-stress 	[12]
3.	Stem	Tinocordifolin 18-norclerodane, glucoside, Furanoid diterpene, glucoside, Tinocordiside, Tinocordifolioside, Cordioside, Cordifolioside, Syringin, Syringinapiosylglycoside, Pregnane glycoside, Palmatosides, Cordifolioside A, B, C, D and E, Glycosides Sesquiterpenoid Berberine, Choline, Tembetarine, Magnoflorine, Tinosporin, Palmetine, Isocolumbin, Aporphine, alkaloids, Jatrorrhizine, Tetrahydropalmatine,	<ul style="list-style-type: none"> ➤ Respiratory tract infections ➤ Skin diseases ➤ Antidote to snake and scorpion sting ➤ Anti-hyperglycemic ➤ Enhance immune response ➤ Anti-carcinogenic ➤ Anti-inflammatory 	[13]
4.	Shoot	β -sitosterol, δ -sitosterol, 20 β -hydroxyecdysone, Ecdysterone, Makisterone A, Giloinsterol Steroids β -sitosterol, δ -sitosterol, 20 β -hydroxyecdysone, Ecdysterone, Makisterone A, Giloinsterol	<ul style="list-style-type: none"> ➤ Anti-carcinogenic ➤ Anti-pyretic 	Singh et al. 2003
5.	Bark	Tinosporofuranol, tinosporafurandioli, tinosporaclerodanol and tinosporaclerodanoid, β - sitosterol	<ul style="list-style-type: none"> ➤ Anti-inflammatory ➤ Antioxidant 	[15]

Table 3. Various chemical constituents, partly used, active components and biological roles of *Tinospora cordifolia* in humans

Sr. no.	Active component	Compound names	Effects in humans	Part used	References	
1.	Alkaloids	➤ Berberine	Anti-cancer, antiviral	Stem, root	[10]	
		➤ Choline				
		➤ Palmatine	infections,			[18]
		➤ Tembetarine	Inflammation,			[19]
		➤ Magnoflorine	and immune-			
		➤ Tetrahydroplamatine	modulatory roles			[20]
		➤ Tinosporin	Neurological,			
		➤ Isocolumbin	Psychiatric			
2.	Glycosides	➤ Jatrorrhizine	Conditions	Stem		
		➤ Aporphine alkaloids	Anti-diabetes			
		➤ 18-norclerodane glucoside	Treats neurological disorders like			[21]
		➤ Furanoid diterpene glucoside	ALS, Parkinsons', dementia, motor			[22]
		➤ Tinocordiside	and cognitive deficits, and neuronloss in spine and			
		➤ Tinocordifolioside	hypothalamus.			
		➤ Cordioside	Immunomodulation: IgG increase			
➤ Palmatosides	and macrophage activation. Inhibits NF-κB and act as nitric oxide scavengers to show anti-cancer activities					
3.	Steroids	➤ β-sitosterol	IgA neuropathy, glucocorticoid	Stems, aerial parts	[23]	
		➤ hydroxy ecdysone	induced osteoporosis in early		[24]	
		➤ Ecdysterone	inflammatory arthritis, induce			
		➤ Giloinsterol	cell cycle arrest in G2/M phase and apoptosis through c-Myc suppression. Inhibits TNF-α, IL-1 β, IL-6 and COX-2. Activates NF-κB			
4.	Aliphatic compounds	➤ Octacosanol	Anti-nociceptive and	Whole plant	[25]	
		➤ Heptacosanol	anti-inflammatory. Protection		[26]	
		➤ Nonacosan-15-one dichloromethane	against 6-hydroxydopamine induced parkinsonism in rats. Down-regulate VEGF and inhibits TNF-α from binding to the DNA			

Sr. no.	Active component	Compound names	Effects in humans	Part used	References
5.	Diterpenoid lactones	<ul style="list-style-type: none"> ➤ Furanolactone ➤ Clerodane derivatives ➤ [(5R,10R)-4R-8R-dihydroxy-2S-3R: ➤ 15,16-diepoxy-cleroda-13 (16), ➤ 14-dieno-17,12S: 18,1S-dilactone] ➤ Tinosporides 	<p>Vasorelaxant: relaxes Norepinephrine induced contractions. Inhibits Ca⁺⁺influx. Anti-inflammatory, anti-microbial, anti-hypertensive, anti-viral. Induce apoptosis in leukemia by activating caspase-3 and bax, inhibits bcl-2</p>	Whole plant	[27] [28]
6.	Others	<ul style="list-style-type: none"> ➤ 3, (a,4-di hydroxy-3-methoxy-benzyl)-4- ➤ (4-hydroxy-3-methoxy-benzyl)-tetrahydrofuran ➤ Jatrorrhizine ➤ N-trans-feruloyl tyramine ➤ Giloin ➤ Tinosporic acid 	<p>Protease inhibitors for HIV and drug resistant HIV. Tyramine is a neuro-modulator. Used to treat anxiety and depression by inactivating neurotransmitters</p>	Root	[29] [30]

Abbreviations: NF- κ B-Nuclear factor-kappa-B, VEGF-Vascular endothelial cell growth factor, TNF-Tumor necrosis factor, IL-Interleukin, COX-Cyclooxygenase, ALS-Amyotrophic, Lateral Sclerosis, IgG-Immunoglobulin G, IgA-Immunoglobulin A

Guduchi is considered one of the best Rasayans and it is unusual potent versatility. Guduchi is known to be a rich source of trace elements (Zinc & Copper) which act as antioxidants & protects cells from the damaging effects of oxygen radicals generated during immune activation. Rasayan effect of Guduchi can be used to heal & prevent infections. Rasayana used as a universal vaccine for any diseases. Rasayana chikitsa is mainly used to maintain the health of individuals although it can be used for diseased also.

Rasa- Tikta, Katu.

Guna- Laghu, Snigha.

Veerya- Ushna.

Vipaka- Madhura

Doshagnata- Tridoshashashamaka

Rogagnata- Kushtha, vatarakta, Netraroga

Karma- Kusthaghna, deepana, Sangrahi, Balya

Prabhava- Tridoshanara, Vishaghna, cure

Rasa- Taste appreciation of the substances by chemical receptors on the tongue, sweet, sour, salt, bitter, pungent and astringent.

Guna- Ten pairs of opposite or mirror image attributes, attribute or property of any substance.

Veerya- Potency, Ushna-hot, Sheeta-cold, Vipakaintestinal digestion and tissue metabolism, Madhuraneutral, Amla-acidic, Katu-alkaline, Prabhava-specification through specialized receptors.

10. MEDICINAL PROPERTIES

- 1. Immunomodulatory activity:** The alcoholic extract of *T. cordifolia* showed significant immunomodulatory effects. At standard dose, extract increased the α -amylase activity and cellularity of bone marrow in rats. It had been observed by some researchers that some active compounds viz; 11-hydroxymustakone, N-methyl-2-pyrrolidone, N-formylannonain, cordifolioside A,

magnoflorine, tinocordiside and syringing showed immunomodulatory activity [31]

- 2. Anti-inflammatory activity:** The water extract of stem part showed anti-inflammatory role in albino rats. It has significantly suppressed acute inflammatory response caused by carrageenin extract when applied orally [15].
- 3. Hepato-suppression:** The extract of the whole plant showed protection against CCl_4 because it causes hepato-cellular changes after forming proteins or by forming bioaction of CCl_4 and accelerated toxification. It also showed potential to reduce the effect of free radicals and antioxidant activity with the suppression of lipid peroxidation, therefore this plant considered as hepatoprotective agent [32].
- 4. Anti-HIV activity:** TCE reduced the recurrent resistance of HIV virus and enhancing the therapeutic outcome [33]. Anti-HIV effects of TCE was revealed a reduction in eosinophil count, stimulation of B lymphocytes, macrophages and polymerphonuclear leucocytes and haemoglobin percentage thus, revealing its promising role of application in the management of the disease [34].
- 5. Anti-cancer activity:** The anti-cancer effects of *T. cordifolia* are mostly studied in animal models. TCE has been shown to have a radioprotective role by significantly increase in body weight, tissue weight, testes-body weight ratio and tubular diameter and inhibit the harmful effects of sub-lethal gamma radiation on testes in male Swiss albino mice [35,36,37,38,39]. In pre-irradiating mice, TCE significantly affected radiation-induced rise in lipid peroxidation and resulted in the decline of GSH concentration in testes. Pre-treatment of HeLa cells by TCE has been shown to decrease the cell viability, increase LDH and decrease in GSH S-transferase activity [40].

11. CONCLUSION

The present study explores the detailed information of *T. cordifolia* and its therapeutic efficiency about the medicinal uses explained in medicinal systems. The phytochemical, pharmaceutical and biological investigation of *T.*

cordifolia reports the versatility and explains its diverse role. It is concluded that this miracle herb had been used traditionally among the various communities across the tribal region of worldwide for ailment of urinary, gastrointestinal, skin, pulmonary, hepatics, gynaecological, inflame-matory and infectious diseases. In addition to this, the species is also well known to treat allergy, tumour and cancer by the traditional and local medicinal practitioners. Almost all parts of the plant are used for curing different but the most frequent part used is rhizome followed by root. In recent times, the old traditional practices are at gradually decline very rapidly and under risk due to rapid modernization hence there is an urgent need for documentation of such tribal species and help to find innovative ways for untap its efficiency used for human welfare in future.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Raghu AV, Geetha SP, Martin G, Balachandran I, Ravindran PN. *In vitro* clonal propagation through mature nodes of *Tinospora cordifolia* (Willd.) Hook F, Thoms. An important ayurvedic medicinal plant. *In vitro cell Dev Biol – Plant*. 2006; 42:584-88.
2. Onkar P, Bangar J, Karodi R. Evaluation of antioxidant activity of traditional formulation *Giloy satva* and hydroalcoholic extract of the *Curculigo orchioides* gaertn). *J App Pharm Sci*. 2012;2(6):209-13.
3. Sangeetha MK, Priya CD, Vasanthi HR. Antidiabetic property of *Tinospora cordifolia* and its active compound is mediated through the expression of Glut-4 in L6 myotubes. *Phytomedicine*. 2013; 20(3-4):246-48.
4. Spandana U, Ali SL, Nirmala T, Santhi M, Sipai Babu SD. A review on *Tinospora cordifolia*. *Int J Curr Pharm Rev Res*. 2013; 4(2):61-8.
5. Singla A. Review of biological activities of "*Tinospora cordifolia*". *Webmed Cent Pharm Sci*. 2010;1(9):WMC0060.
6. Sinha K, Mishr NP, Singh J, Khanuja SPS. *Tinospora cordifolia*, a reservoir plant for therapeutic application. *Indian J Tradit Know*. 2004;3(3):257-70.
7. Shetty BV, Singh V. *Flora of Rajasthan*. 1st edition, Merrut publishers and Distributors, Merrut. 2010;1:756-1000.
8. Singh J, Sinha K, Sharma A, Mishra NP, Khanuja SP. Traditional uses of *Tinospora cordifolia* (Guduchi). *J Med Aromat Plant Sci*. 2003;25:748–51.
9. Hossain MM, Hasan SMR, Akter R, Islam MN, Rashid MJ, Saha MR, et al. Evaluation of analgesic and neuropharmacological properties of the aerial part of *Tinospora cordifolia* Miers. in mice. *S J Pharm Sci*. 2009; 2(2):31-7.
10. Upadhyay AK, Kumar K, Kumar A, Mishra SH. *Tinospora cordifolia* (Wild.) Hook. f. and Thoms. (Guduchi)-validation of the Ayurvedic pharmacology through experimental and clinical studies. *Int J Ayurveda Res*. 2010;1(2): 112–21.
11. Kavitha BT, Shruthi SD, Rai SP, Ramachandra YL. Phytochemical analysis and hepatoprotective properties of *Tinospora cordifolia* against carbon tetrachloride induced hepatic damage in rats. *J Basic Clin Pharm*. 2011;2(3):139-42.
12. Dikshit V, Damre AS, Kulkarni KR, Gokhale A, Saraf MN. Preliminary screening of immunocin for immunomodulatory activity. *Indian J Pharm Sci*. 2000;62:257.
13. Desai RV, Kamat JB, Sainis KB. An immunomodulator from *Tinospora cordifolia* with antioxidant activity in cell-free systems. *J Chem Sci*. 2012;114(6): 713-19.
14. Singh SS, Pandey SC, Srivastava S, Gupta VS, Patro B, Ghosh AC. Chemistry and medicinal properties of *Tinospora cordifolia* (Guduchi). *Indian J Pharm*. 2003; 35:83-91.
15. Pendse VK, Dadhich AP, Mathur PN, Madam MSB. Anti-inflammatory, immunosuppressive and some related pharmacological actions of the water extract of Neem Giloe (*Tinospora cordifolia*): A preliminary report. *Indian J Pharm*. 1977;9:221-24.
16. Neeraja PV, Margaret E. Amruthavalli (*Tinospora cordifolia*) multipurpose rejuvenator. *Int J Pharm Biol Chem Sci*. 2013;3(2):233-41.

17. Sharma PC, Yelne MB, Dennis TJ. Database on medicinal plants used in Ayurveda. Central council of research in Ayurveda & Siddha, New Delhi; 2005.
18. Rout GR. Identification of *Tinospora cordifolia* (Willd.) Miers ex Hook F & Thomas using RAPD markers. Z Naturf Sch C. 2006;61:118-22.
19. Patel SS, Shah RS, Goyal RK. Antihyperglycemic, antihyperlipidemic and antioxidant effects of Dihar, a polyherbal ayurvedic formulation in streptozotocin induced diabetic rats. Indian J Exp Biol. 2009;47:564-70.
20. Patel MB, Mishra S. Hypoglycemic activity of alkaloidal fraction of *Tinospora cordifolia*. Phytomedicine. 2011;18:1045-52.
21. Ly PT, Singh S, Shaw CA. Novel environmental toxins: Steryl glycosides as a potential etiological factor for age-related neurodegenerative diseases. J Neurosci Res. 2007;85:231-7.
22. Sengupta S, Mukherjee A, Goswami R, Basu S. Hypoglycemic activity of the antioxidant saponarin, characterized as alpha-glucosidase inhibitor present in *Tinospora cordifolia*. J Enzyme Inhib Med Chem. 2009;24:684-90.
23. Lv J, Xu D, Perkovic V, Ma X, Johnson DW, Woodward M, et al. Corticosteroid therapy in IgA nephropathy. J Am Soc Nephrol. 2012;23:1108-16.
24. Lee IA, Kim EJ, Kim DH. Inhibitory effect of β -sitosterol on TNBS-induced Colitis in mice. Planta Med. 2012;78: 896-8.
25. De Oliveira AM, Conserva LM, de Souza Ferro JN, de Almeida Brito F, Lyra Lemos RP, Barreto E. Antinociceptive and anti-inflammatory effects of octacosanol from the leaves of *Sabicea grisea* var. *Grisea* in mice. Int J Mol Sci. 2012;13: 1598-1611.
26. Taylor JC, Rapport L, Lockwood GB. Octacosanol in human health. Nutrition. 2003;19:192-5.
27. Sriramaneni RN, Omar AZ, Ibrahim SM, Amirin S, Mohd Zaini A Vasorelaxant effect of diterpenoid lactones from *Andrographis paniculata* chloroform extract on rat aortic rings. Pharmacognosy Res. 2010;2:242-6.
28. Dhanasekaran M, Baskar AA, Ignacimuthu S, Agastian P, Duraipandiyan V. Chemopreventive potential of Epoxy clerodane diterpene from *Tinospora cordifolia* against diethylnitrosamine- induced hepatocellular carcinoma. Invest New Drugs. 2009;27: 347-55.
29. Ghosh AK, Martyr CD, Steffey M, Wang YF, Agniswamy J, Amano M, et al. Design of substituted bis - Tetrahydrofuran (bis-THF) - derived potent HIV - 1 protease inhibitors, protein-ligand X-ray structure, and convenient syntheses of bis - THF and Substituted bis - THF Ligands. ACS Med Chem Lett. 2011;2:298-302.
30. Mukherjee R, De UK, Ram GC. Evaluation of mammary gland immunity and therapeutic potential of *Tinospora cordifolia* against bovine subclinical mastitis. Trop Anim Health Prod. 2010;42:645-51.
31. Upadhyaya R, PR, Sharma V, Anita KV. Assessment of the multifaceted immunomodulatory potential of the aqueous extract of *Tinospora cordifolia*. Res J Chem Sci. 2011;1:71-9.
32. Sharma V, Pandey D. Beneficial effects of *Tinospora cordifolia* on blood profiles in male mice exposed to lead. Toxicol Int. 2010;17:8-11.
33. Kalikar MV, Thawani VR, Varadpande UK, Sontakke SD, Singh RP, Khyani RK. Immunomodulatory effect of *Tinospora cordifolia* extract in human immunodeficiency virus positive patients. Indian J Pharmacol. 2008;40:107-10.
34. Akhtar S. Use of *Tinospora cordifolia* in HIV infection. Indian J Pharmacol. 2010: 42:57.
35. Rao SK, Rao PS. Alteration in the radiosensitivity of HeLa cells by dichloromethane extract of guduchi (*Tinospora cordifolia*). Integr Cancer Ther. 2010;9:378-84.
36. GawhareVikesh S. A review on guduchi through Ayurvedic texts. International Ayurvedic Medical Journal. 2013;1(3):1-7.
37. Pratibha Baghel. Plant of versatile properties: A review of *Tinospora Cordifolia* (Guduchi). International Journal of Agriculture Innovations and Research. 2017;5(5):751-753.

38. Saha S, Ghosh S. *Tinospora cordifolia*: One plant, many roles. *Ancient Science of Life*. 2012;31(4):151.
39. Sevindik M, Akgul H, Pehlivan M, Selamoglu Z. Determination of therapeutic potential of *Mentha longifolia* ssp. *longifolia*. *Fresen Environ Bull*. 2017;26: 4757-4763.
40. Pehlivan M, Mohammed FS, Sevindik M, Akgul H. Antioxidant and oxidant potential of *Rosa canina*. *Eurasian Journal of Forest Science*. 2018;6(4):22-25.

© 2019 Antul et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/49114>